

CONFORMAL MEASURE RIGIDITY AND ERGODICITY OF HOROSPHERICAL FOLIATIONS

DONGRYUL M. KIM

ABSTRACT. In this paper, we prove two main theorems: conformal measure rigidity and ergodicity of horospherical foliations, especially in higher rank. Both theorems are new even for relatively Anosov groups.

First, we establish a higher rank extension of rigidity theorems of Sullivan, Tukia, Yue, and Kim-Oh for representations of rank one discrete subgroups of divergence type, in terms of the push-forwards of conformal measures via boundary maps. We consider a certain class of higher rank discrete subgroups, which we call hypertransverse subgroups. It includes all rank one discrete subgroups, Anosov subgroups, relatively Anosov subgroups, and their subgroups. Our proof of the rigidity theorem generalizes the idea of Kim-Oh to self-joinings of higher rank hypertransverse subgroups, overcoming the lack of $CAT(-1)$ geometry on symmetric spaces. In contrast to the work of Sullivan, Tukia, and Yue, our argument is closely related to the study of horospherical foliations.

We also show the ergodicity of horospherical foliations with respect to Burger-Roblin measures. This generalizes the classical work of Hedlund, Burger, and Roblin in rank one and of Lee-Oh for Borel Anosov subgroups in higher rank. Moreover, we describe the ergodic decomposition of Burger-Roblin measures and Bowen-Margulis-Sullivan measures when a given parabolic subgroup is minimal.

CONTENTS

1. Introduction	2
2. Preliminaries	10
3. Busemann maps, conformal measures and essential subgroups	14
4. Graph-conformal measures for self-joinings	16
5. Myrberg limit sets	19
6. δ -hyperbolic spaces	24
7. Essential subgroups for graph-conformal measures	36
8. Singularity of the graph-conformal measure	43
9. Deformations of transverse representations	44
10. Horospherical foliations and Burger-Roblin measures	45
References	48

1. INTRODUCTION

The celebrated rigidity theorem of Mostow [33] (see also Prasad [34, Theorem B] for non-uniform lattices) states that if Γ is a lattice of $G = \text{Isom}^+(\mathbb{H}^n)$, $n \geq 3$, then any discrete faithful representation $\rho : \Gamma \rightarrow G$ extends to a Lie group isomorphism $G \rightarrow G$. The crucial part of Mostow's proof is that there exists a ρ -equivariant homeomorphism $f : \mathbb{S}^{n-1} \rightarrow \mathbb{S}^{n-1}$ which is quasiconformal.

Sullivan showed a rigidity theorem for discrete faithful representations of a general discrete subgroup of G , extending Mostow's rigidity theorem. Let $\Gamma < G$ be a discrete subgroup and denote its limit set by $\Lambda_\Gamma \subset \mathbb{S}^{n-1}$. We also denote the critical exponent of Γ by δ_Γ , which is defined as the abscissa of convergence of the Poincaré series $\sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)}$, $o \in \mathbb{H}^n$. We say that Γ is of divergence type if the Poincaré series diverges at $s = \delta_\Gamma$. In this case, there exists a unique δ_Γ -dimensional Γ -conformal measure ν_Γ on \mathbb{S}^{n-1} and it charges the full mass on the conical limit set, in particular $\nu_\Gamma(\Lambda_\Gamma) = 1$ [39]. Sullivan's conformal measure rigidity theorem is as follows:

Theorem 1.1 (Sullivan, [40, Theorem 5]). *Let $\Gamma < G$ be a discrete subgroup of divergence type and $\rho : \Gamma \rightarrow G$ a discrete faithful representation such that $\delta_{\rho(\Gamma)} = \delta_\Gamma$. If ρ admits an equivariant continuous embedding $f : \Lambda_\Gamma \rightarrow \mathbb{S}^{n-1}$ and¹*

$$\nu_{\rho(\Gamma)} \ll f_* \nu_\Gamma$$

for some $\delta_{\rho(\Gamma)}$ -dimensional $\rho(\Gamma)$ -conformal measure $\nu_{\rho(\Gamma)}$, then ρ extends to a Lie group isomorphism $G \rightarrow G$.

Later, Tukia [41, Theorem 3C] showed that the condition $\delta_{\rho(\Gamma)} = \delta_\Gamma$ is not necessary in Theorem 1.1. Since a quasiconformal homeomorphism preserves the Lebesgue measure class on \mathbb{S}^{n-1} , Mostow's rigidity theorem also follows from Theorem 1.1. Push-forwards of conformal measures via boundary maps were also used in the rigidity theorem of Besson-Courtois-Gallot ([4], [5]).

Sullivan and Tukia's proofs use the ergodicity of the geodesic flow with respect to the Bowen-Margulis-Sullivan measure on the unit tangent bundle of $\Gamma \backslash \mathbb{H}^n$ to deduce the conformality of the boundary map f relying on the negatively curved geometry of the real hyperbolic space. Generalizing this idea, Yue extended Theorem 1.1 to general rank one spaces [43, Theorem A].

In our recent series of work with Oh ([23], [22], [24]), we introduced a new approach in the study of rigidity problems on a representation ρ of a discrete subgroup (possibly with infinite-covolume), that is, considering the self-joining of Γ via ρ and relating to the higher rank dynamics of the self-joining subgroup. Especially in [22], the conformal measure rigidity was studied for representations of a rank one discrete subgroup into a simple real algebraic group of general rank, whose orbit maps are extended to

¹The notation $\nu_1 \ll \nu_2$ means that ν_1 is absolutely continuous with respect to ν_2 .

Furstenberg boundaries. It recovers Theorem 1.1 as well as the work of Tukia and Yue.

Conformal measure rigidity in general rank. In this paper, we establish the conformal measure rigidity theorem for representations of discrete subgroups of a general-rank simple real algebraic group. We consider a certain class of discrete subgroups, which we call hypertransverse subgroups. This includes rank one discrete subgroups, Anosov subgroups, relatively Anosov subgroups, and their subgroups.

We first introduce some terminologies and notations that we use throughout the paper. Let G be a connected semisimple real algebraic group. Let $P < G$ be a minimal parabolic subgroup with a fixed Langlands decomposition $P = MAN$ where A is a maximal real split torus of G , M is the maximal compact subgroup of P commuting with A and N is the unipotent radical of P . Let $\mathfrak{g} = \text{Lie } G$ and $\mathfrak{a} = \text{Lie } A$. Fix a positive Weyl chamber $\mathfrak{a}^+ < \mathfrak{a}$ and set $A^+ = \exp \mathfrak{a}^+$. We fix a maximal compact subgroup $K < G$ such that the Cartan decomposition $G = KA^+K$ holds. We use the notation $\mu : G \rightarrow \mathfrak{a}^+$ for the Cartan projection, defined by the condition that $g \in K \exp \mu(g)K$ for $g \in G$. We have the associated Riemannian symmetric space $X = G/K$ and write $o = [K] \in X$.

Let Π denote the set of all simple roots for $(\mathfrak{g}, \mathfrak{a}^+)$. As usual, the Weyl group is the quotient of the normalizer of A in K by the centralizer of A in K . We also denote the opposition involution by $i : \mathfrak{a} \rightarrow \mathfrak{a}$. It induces an involution on Π which we denote by the same notation i . Throughout the paper, we fix a non-empty subset

$$\theta \subset \Pi.$$

Let $\mathfrak{a}_\theta = \bigcap_{\alpha \in \Pi - \theta} \ker \alpha$ and let $p_\theta : \mathfrak{a} \rightarrow \mathfrak{a}_\theta$ be the unique projection, invariant under all Weyl elements fixing \mathfrak{a}_θ pointwise. We write $\mu_\theta := p_\theta \circ \mu$. Let P_θ be the standard parabolic subgroup corresponding to θ (our convention is that $P = P_\Pi$) and consider the θ -boundary:

$$\mathcal{F}_\theta = G/P_\theta.$$

We say that $\xi \in \mathcal{F}_\theta$ and $\eta \in \mathcal{F}_{i(\theta)}$ are in general position if the pair (ξ, η) belongs to the unique open G -orbit in $\mathcal{F}_\theta \times \mathcal{F}_{i(\theta)}$ under the diagonal action of G .

Conformal measures. Denote by $\mathfrak{a}_\theta^* = \text{Hom}(\mathfrak{a}_\theta, \mathbb{R})$ the space of all linear forms on \mathfrak{a}_θ . For $\psi \in \mathfrak{a}_\theta^*$ and a closed subgroup $\Delta < G$, a Borel probability measure ν on \mathcal{F}_θ is called a (Δ, ψ) -conformal measure (with respect to $o \in X$) if

$$\frac{dg_*\nu}{d\nu}(\xi) = e^{\psi(\beta_\xi^\theta(o, go))} \quad \text{for all } g \in \Delta \text{ and } \xi \in \mathcal{F}_\theta$$

where $g_*\nu(D) = \nu(g^{-1}D)$ for any Borel subset $D \subset \mathcal{F}_\theta$ and β_ξ^θ denotes the \mathfrak{a}_θ -valued Busemann map (Definition 3.1). By a Δ -conformal measure, we

mean a (Δ, ψ) -conformal measure for some $\psi \in \mathfrak{a}_\theta^*$. The linear form ψ plays a role of *dimension* of ν .

We say that a (Δ, ψ) -conformal measure ν on \mathcal{F}_θ is of *divergence type* if ψ is (Δ, θ) -proper² and $\sum_{g \in \Delta} e^{-\psi(\mu_\theta(g))} = \infty$. The (Δ, θ) -properness hypothesis guarantees that the abscissa of convergence of the Poincaré series $\sum_{g \in \Delta} e^{-s\psi(\mu_\theta(g))}$, which we denote by δ_ψ , is well-defined.

Hypertransverse subgroups. Let $\Gamma < G$ be a Zariski dense discrete subgroup. We denote by $\Lambda^\theta := \Lambda_\Gamma^\theta \subset \mathcal{F}_\theta$ the limit set of Γ in \mathcal{F}_θ , which is the unique Γ -minimal subset of \mathcal{F}_θ (Definition 2.8). A discrete subgroup Γ is called θ -transverse if

- Γ is θ -regular, i.e., $\liminf_{\gamma \in \Gamma} \alpha(\mu_\theta(\gamma)) = \infty$ for all $\alpha \in \theta$; and
- Γ is θ -antipodal, i.e., any two distinct $\xi, \eta \in \Lambda^{\theta \cup i(\theta)}$ are in general position.

Most of the known examples of transverse subgroups come with nice actions on Gromov hyperbolic spaces. In this regard, we consider the following subclass:

Definition 1.2. A θ -transverse subgroup $\Gamma < G$ is called θ -hypertransverse if there exists a proper geodesic Gromov hyperbolic space Z such that

- Γ acts on Z properly discontinuously by isometries;
- there exists a Γ -equivariant homeomorphism

$$\Lambda^Z \rightarrow \Lambda^\theta$$

where Λ^Z is the limit set of Γ in the Gromov boundary ∂Z .

Example 1.3. As mentioned before, any subgroup of an Anosov or a relatively Anosov group (Definition 1.7) is hypertransverse. Indeed, when Γ is a subgroup of an Anosov group Γ_0 , we can take Z to be the Cayley graph of Γ_0 . For a subgroup Γ of a relatively Anosov group Γ_0 , we can set Z to be the Groves-Manning cusp space of Γ_0 .

It seems that most transverse subgroups are hypertransverse. We do not know of an example of a transverse subgroup which is not hypertransverse.

Rigidity theorems. Let G_1, G_2 be connected simple real algebraic groups. Let θ_1 and θ_2 be non-empty subsets of the set of simple roots of G_1 and G_2 respectively. Here is our main rigidity theorem:

Theorem 1.4 (Conformal measure rigidity). *Let $\Gamma < G_1$ be a Zariski dense θ_1 -hypertransverse subgroup. Let $\rho : \Gamma \rightarrow G_2$ be a Zariski dense θ_2 -regular³ faithful representation with a pair of ρ -equivariant continuous embeddings*

²a linear form $\psi \in \mathfrak{a}_\theta^*$ is called (Δ, θ) -proper if $\psi \circ \mu_\theta : \Delta \rightarrow [-\varepsilon, \infty)$ is a proper map for some $\varepsilon > 0$.

³i.e. $\rho(\Gamma)$ is θ_2 -regular

$f : \Lambda^{\theta_1} \rightarrow \mathcal{F}_{\theta_2}$ and $f_i : \Lambda^{i(\theta_1)} \rightarrow \mathcal{F}_i(\theta_2)$. If there exists a Γ -conformal measure ν_Γ of divergence type such that

$$\nu_{\rho(\Gamma)} \ll f_*\nu_\Gamma$$

for some $\rho(\Gamma)$ -conformal measure $\nu_{\rho(\Gamma)}$, then ρ extends to a Lie group isomorphism $G_1 \rightarrow G_2$.

Remark 1.5. We emphasize that there is no additional assumption on the conformal measure $\nu_{\rho(\Gamma)}$ and its associated linear form, such as $(\rho(\Gamma), \theta_2)$ -properness. Moreover, we do not assume that the image $\rho(\Gamma)$ is transverse.

We note that if a ρ -equivariant map $\Lambda^{\theta_1} \rightarrow \mathcal{F}_{\theta_2}$ exists, then it is unique (Lemma 4.3). We also note that if $\psi \in \mathfrak{a}_{\theta_1}^*$ is (Γ, θ_1) -proper and the associated Poincaré series diverges at δ_ψ , then there exists a unique $(\Gamma, \delta_\psi, \psi)$ -conformal measure on \mathcal{F}_{θ_1} and has support Λ^{θ_1} ([13], [26]).

When $\text{rank } G_1 = 1$, a Γ -conformal measure ν is of divergence type if and only if Γ is of divergence type and ν is the δ_Γ -dimensional Γ -conformal measure. Hence Theorem 1.4 generalizes Theorem 1.1 as well as the work of Tukia [41] and Yue [43] in the case that $\text{rank } G_2 = 1$, and the work of Kim-Oh [22] for general G_2 .

We also show the following:

Theorem 1.6 (Singularity between conformal measures). *Let G be a semisimple real algebraic group and $\Gamma < G$ be a Zariski dense θ -hypertransverse subgroup. Let ν be a (Γ, ψ) -conformal measure of divergence type. Then for any (Γ, ψ') -conformal measure ν' with $\psi' \neq \psi$, we have*

$$\nu' \not\ll \nu.$$

In particular, if ν' is further assumed to be of divergence type, then

$$\nu' \perp \nu.$$

Theorem 1.6 also follows from the work of Lee-Oh [32] and of Sambarino [37] when Γ is θ -Anosov. Recently, Blayac-Canary-Zhu-Zimmer [6] showed a related result on singularity/absolute continuity between two Patterson-Sullivan measures in a more abstract setting, assuming that they are supported on the limit set.

Relatively Anosov subgroups. Theorem 1.4 and Theorem 1.6 apply to any Zariski dense subgroup Γ of a relatively Anosov subgroup Δ (see Example 1.3). The notion of relatively Anosov subgroups (resp. Anosov subgroups) was introduced as a higher rank extension of geometrically finite subgroups (resp. convex cocompact subgroups); see ([28], [17], [21], [16], [12], [44]).

Definition 1.7. Let $\Delta < G$ be a θ -transverse subgroup and \mathcal{P} a finite collection of subgroups in Γ . We say that $\Delta < G$ is θ -Anosov relative to \mathcal{P} if Δ is a hyperbolic group relative to \mathcal{P} , and there exists a Δ -equivariant homeomorphism from the Bowditch boundary $\partial(\Delta, \mathcal{P})$ to Λ_Δ^θ . When $\mathcal{P} = \emptyset$, Δ is called θ -Anosov.

Rigidity of transverse representations. We also consider a conjugate between two transverse representations, which can be regarded as a deformation between them. Let (Z, d_Z) be a proper geodesic Gromov hyperbolic space. Let $\Delta < \text{Isom}(Z)$ be a subgroup of isometries of Z acting properly discontinuously on Z . We denote by $\Lambda_\Delta^Z \subset \partial Z$ its limit set in the Gromov boundary.

For $i = 1, 2$, let G_i be a simple real algebraic group and $\rho_i : \Delta \rightarrow G_i$ a Zariski dense θ_i -transverse representation, in the sense that $\rho_i(\Delta) < G_i$ is a θ_i -transverse subgroup and that there exists a ρ_i -equivariant homeomorphism $f_i : \Lambda_\Delta^Z \rightarrow \Lambda_{\rho_i(\Delta)}^{\theta_i}$. We set $\Gamma_i := \rho_i(\Delta)$. Then the isomorphism $\rho := \rho_2 \circ \rho_1|_{\Gamma_1}^{-1}$ conjugates two representations ρ_1 and ρ_2 :

$$\begin{array}{ccc} & & \Gamma_1 \\ & \nearrow^{\rho_1} & \downarrow \rho \\ \Delta & & \\ & \searrow_{\rho_2} & \Gamma_2 \end{array}$$

The following is an immediate consequence of Theorem 1.4 where f is the ρ -boundary map:

Theorem 1.8. *Let $\rho : \Gamma_1 \rightarrow \Gamma_2$ be a conjugate between two Zariski dense transverse representations $\rho_1 : \Delta \rightarrow \Gamma_1$ and $\rho_2 : \Delta \rightarrow \Gamma_2$. Suppose that ρ does not extend to a Lie group isomorphism $G_1 \rightarrow G_2$. For any Γ_1 -conformal measure ν_1 of divergence type and Γ_2 -conformal measure ν_2 ,*

$$\nu_2 \not\ll f_*\nu_1.$$

In particular, if ν_2 is further assumed to be of divergence type, then

$$\nu_2 \perp f_*\nu_1.$$

In other words, we have the following rigidity theorem for transverse representations:

Theorem 1.9. *If there exist a Γ_1 -conformal measure ν_1 of divergence type such that*

$$\nu_2 \ll f_*\nu_1$$

for some Γ_2 -conformal measure ν_2 , then ρ extends to a Lie group isomorphism $G_1 \rightarrow G_2$.

Ergodicity of horospherical foliations. The horospherical foliation \mathcal{H} of the unit tangent bundle $T^1(\mathbb{H}^2)$ is a collection of horospheres stable under the geodesic flow. This can be identified as follows:

$$\mathcal{H} = \partial\mathbb{H}^2 \times \mathbb{R} = \text{PSL}_2(\mathbb{R})/N$$

where $N = \left\{ \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} : s \in \mathbb{R} \right\}$. Hedlund showed that when Γ is a lattice in $\text{PSL}_2(\mathbb{R})$, the Γ -action on \mathcal{H} is ergodic with respect to the Haar measure

[19]. Later, Burger [10] proved that any convex cocompact $\Gamma < \mathrm{PSL}_2(\mathbb{R})$ with $\delta_\Gamma > 1/2$ acts ergodically on \mathcal{H} with respect to the measure

$$e^{\delta_\Gamma t} d\nu_\Gamma dt$$

where $d\nu_\Gamma$ is the unique δ_Γ -dimensional Γ -conformal measure on $\partial\mathbb{H}^2$ and dt is a Lebesgue measure on \mathbb{R} . This measure on $\mathcal{H} = \partial\mathbb{H}^2 \times \mathbb{R}$ is called the Burger-Roblin measure which we denote by m_Γ^{BR} . Roblin extended these results to a more general setting:

Theorem 1.10. [36, Corollary 2.3] *Let (X, d) be a proper $\mathrm{CAT}(-1)$ -space and $\Gamma < \mathrm{Isom}(X)$ a discrete subgroup. Suppose that $\sum_{\gamma \in \Gamma} e^{-\delta_\Gamma d(o, \gamma o)} = \infty$ and the length spectrum of Γ is non-arithmetic⁴. Then the Γ -action on the horospherical foliation of X with respect to m_Γ^{BR} is ergodic.*

When X is a rank one symmetric space, the length spectrum of any non-elementary discrete subgroup $\Gamma < \mathrm{Isom}(X)$ is non-arithmetic [27]. Hence Theorem 1.10 implies that any discrete subgroup $\Gamma < \mathrm{Isom}(X)$ of divergence type acts ergodically on the horospherical foliation of $T^1(X)$ with respect to the Burger-Roblin measure.

We extend Theorem 1.10 to higher rank settings. In the rest of the introduction, let G be a connected semisimple real algebraic group and fix a non-empty $\theta \subset \Pi$. We then have the Langlands decomposition $P_\theta = N_\theta S_\theta A_\theta$ where $A_\theta = \exp \mathfrak{a}_\theta$, S_θ is an almost direct product of a semisimple algebraic group and a compact central torus, and N_θ is the unipotent radical of P_θ . The θ -horospherical foliation is defined as

$$\mathcal{H}_\theta := \mathcal{F}_\theta \times \mathfrak{a}_\theta = G/N_\theta S_\theta$$

analogous to the rank one setting. Since A_θ normalizes $N_\theta S_\theta$, it acts on \mathcal{H}_θ on the right by multiplication (see (10.1)).

Let $\Gamma < G$ be a discrete subgroup and ν a (Γ, ψ) -conformal measure on \mathcal{F}_θ . The Burger-Roblin measure associated to ν is a Γ -invariant Radon measure on \mathcal{H}_θ defined as

$$dm_\nu^{\mathrm{BR}}(\xi, u) := e^{\psi(u)} d\nu(\xi) du$$

where du is the Lebesgue measure on \mathfrak{a}_θ . Note that

$$\mathrm{supp} m_\nu^{\mathrm{BR}} = \{(\xi, u) \in \mathcal{H}_\theta : \xi \in \mathrm{supp} \nu\}.$$

Here is a higher rank version of Theorem 1.10 relating the ergodicity of $(\mathcal{H}_\theta, \Gamma, \nu_\nu^{\mathrm{BR}})$ with the divergence of the ψ -Poincaré series:

Theorem 1.11. *Let $\Gamma < G$ be a Zariski dense θ -hypertransverse subgroup. For any Γ -conformal measure ν of divergence type,*

the Γ -action on the horospherical foliation $(\mathcal{H}_\theta, m_\nu^{\mathrm{BR}})$ is ergodic.

⁴The non-arithmeticity of the length spectrum means that the set of all lengths of closed geodesics in $\Gamma \backslash X$ generates a dense subgroup of $(\mathbb{R}, +)$.

Horospherical actions on $\Gamma \backslash G$. Considering the case $\theta = \Pi$, we have $N_\Pi = N$ and $S_\Pi = M$, and hence

$$\mathcal{H}_\Pi = G/NM.$$

By the Iwasawa decomposition $G = KAN = KP$, the Furstenberg boundary \mathcal{F} is identified with K/M . For a (Γ, ψ) -conformal measure ν on \mathcal{F} , we denote by $\hat{\nu}$ the M -invariant lift of ν to K . We then define the following Γ -invariant measure on G : for $g = k(\exp u)n \in KAN$,

$$(1.1) \quad d\hat{m}_\nu^{\text{BR}}(g) := e^{\psi(u)} d\hat{\nu}(k) dudn$$

where dn is the Haar measure on N . The measure \hat{m}_ν^{BR} is the NM -invariant lift of m_ν^{BR} to G , and induces the measure on $\Gamma \backslash G$ which we also call the Burger-Roblin measure and denote by \hat{m}_ν^{BR} , abusing notations.

We consider the horospherical action on $(\Gamma \backslash G, \hat{m}_\nu^{\text{BR}})$, given as the right multiplication by NM . Since any conformal measure of divergence type is supported on the limit set [26, Theorem 1.5], Theorem 1.11 can be rephrased as follows in this case:

Theorem 1.12. *Let $\Gamma < G$ be a Zariski dense Π -hypertransverse subgroup. For any Γ -conformal measure ν on \mathcal{F} of divergence type,*

$$\text{the } NM\text{-action on } (\Gamma \backslash G, \hat{m}_\nu^{\text{BR}}) \text{ is ergodic.}$$

In particular, for \hat{m}_ν^{BR} -a.e. $x \in \Gamma \backslash G$, we have

$$\overline{xNM} = \{[g] \in \Gamma \backslash G : gP \in \Lambda^\Pi\}.$$

Theorem 1.12 applies to the images of cusped Hitchin representations, which are well-known examples of relatively Π -Anosov subgroups (see [8] and [11]).

Theorem 1.11 and Theorem 1.12 extend the work of Lee-Oh [32] on Π -Anosov subgroups. See also [29] for a certain unique ergodicity result of Burger-Roblin measures for special types of Π -Anosov subgroups.

Remark 1.13. Lee-Oh described the ergodic decomposition of the Burger-Roblin and Bowen-Margulis-Sullivan measures on $\Gamma \backslash G$ for a Zariski dense Π -Anosov subgroup $\Gamma < G$ [31]. In view of Theorem 1.12, their argument applies to Π -hypertransverse subgroups, and hence yields similar ergodic decomposition theorems. See Theorem 10.4 for a precise statement.

On the proofs. As mentioned before, Theorem 1.4 was proved by Kim-Oh [22] when Γ is a rank one discrete subgroup and ρ is a certain representation into a higher rank simple group. The major point of this paper is to extend the proof of [22] to higher rank. Under the rank one assumption on Γ , the symmetric space is $\text{CAT}(-1)$, and hence Busemann functions and Gromov products behave nicely enough. However, when Γ is of higher rank, the symmetric space is neither negatively curved nor $\text{CAT}(-1)$, and hence it requires additional ideas to prove Theorem 1.4 in this generality. Even under the hypertransverse hypothesis on Γ , it only admits an action on a

Gromov hyperbolic space, and the coarse nature of the geometry of Gromov hyperbolic spaces still presents several non-trivial difficulties in extending previous works on the conformal measure rigidity [22] and the ergodicity of horospherical foliations ([36], [32]).

In contrast to the proof of Sullivan, Tukia, and Yue, we consider the self-joining of Γ via $\rho : \Gamma \rightarrow G_2$ to prove Theorem 1.4:

$$\Gamma_\rho := (\text{id} \times \rho)(\Gamma) = \{(\gamma, \rho(\gamma)) : \gamma \in \Gamma\},$$

a discrete subgroup of $G := G_1 \times G_2$. As G_1 and G_2 are simple, ρ extends to a Lie group isomorphism $G_1 \rightarrow G_2$ if and only if Γ_ρ is not Zariski dense (Lemma 4.2). Hence we translate the rigidity question on ρ to the Zariski density question on the self-joining Γ_ρ . The idea relating dynamics of self-joinings to rigidity problems in terms of boundary maps originates from the work of Kim-Oh ([23], [24], [22]).

As in [22], we let $A := A_1 \times A_2$ and $A^+ := A_1^+ \times A_2^+$ so that $\mathfrak{a} := \text{Lie } A = \mathfrak{a}_1 \oplus \mathfrak{a}_2$ and $\mathfrak{a}^+ = \mathfrak{a}_1^+ \oplus \mathfrak{a}_2^+$. Denote by $\Pi = \Pi_1 \cup \Pi_2$ the set of all simple roots for $(\text{Lie } G, \mathfrak{a}^+)$ and $\theta = \theta_1 \cup \theta_2$. We have the θ -boundary $\mathcal{F}_\theta = \mathcal{F}_{\theta_1} \times \mathcal{F}_{\theta_2}$ and $\mathfrak{a}_\theta = \mathfrak{a}_{\theta_1} \oplus \mathfrak{a}_{\theta_2}$.

For a (Γ, ψ) -conformal measure ν of divergence type for $\psi \in \mathfrak{a}_{\theta_1}^*$, recall the graph-conformal measure defined by Kim-Oh [22] which is the push-forward

$$\nu_\rho := (\text{id} \times f)_*\nu.$$

As observed in [22], ν_ρ is $(\Gamma_\rho, \sigma_\psi)$ -conformal where $\sigma_\psi \in \mathfrak{a}_\theta^*$ is defined by $\sigma_\psi(u_1, u_2) = \psi(u_1)$ for $(u_1, u_2) \in \mathfrak{a}_{\theta_1} \oplus \mathfrak{a}_{\theta_2}$.

The main technical ingredient of the proof is to show that if Γ_ρ is Zariski dense, the essential subgroup $\mathbb{E}_{\nu_\rho}^\theta(\Gamma_\rho)$ is the whole \mathfrak{a}_θ (Theorem 7.1). The essential subgroup $\mathbb{E}_{\nu_\rho}^\theta(\Gamma_\rho) \subset \mathfrak{a}_\theta$ is defined as the set of $u \in \mathfrak{a}_\theta$ such that for any $\varepsilon > 0$ and a Borel subset $B \subset \mathcal{F}_\theta$ with $\nu_\rho(B) > 0$, the subset

$$B \cap \gamma B \cap \{\xi \in \mathcal{F}_\theta : \|\beta_\xi^\theta(e, \gamma) - u\| < \varepsilon\}$$

has a positive ν_ρ -measure for some $\gamma \in \Gamma_\rho$. That $\mathbb{E}_{\nu_\rho}^\theta(\Gamma_\rho) = \mathfrak{a}_\theta$ implies the singularity of ν_ρ among conformal measures of Γ_ρ , and hence the singularity of $f_*\nu$ between $\rho(\Gamma)$ -conformal measures (Proposition 3.8). Therefore, the non-singularity between $f_*\nu$ and a $\rho(\Gamma)$ -conformal measure forbids Γ_ρ from being Zariski dense and hence ρ extends to a Lie group isomorphism $G_1 \rightarrow G_2$.

To show $\mathbb{E}_{\nu_\rho}^\theta(\Gamma_\rho) = \mathfrak{a}_\theta$, we first prove that the Myrberg limit sets of Γ and of the self-joining Γ_ρ have full ν and ν_ρ -measures respectively (Theorem 5.3), only assuming that Γ is a transverse subgroup. This is based on the ergodicity of an appropriate one-dimensional flow obtained in [26, Theorem 10.2] and is new even for a transverse subgroup Γ .

We then deduce $\mathbb{E}_{\nu_\rho}^\theta(\Gamma_\rho) = \mathfrak{a}_\theta$ from the full ν_ρ -mass of the Myrberg limit set of Γ_ρ under the additional hypothesis that Γ is hypertransverse. The idea of this deduction is influenced by Roblin [36] in the CAT(-1) setting,

by Kim-Oh [22] for self-joinings of rank one discrete subgroups, and by Lee-Oh [32] dealing with Anosov subgroups with respect to minimal parabolic subgroups. The fact that the visual metrics behave nicely in the $\text{CAT}(-1)$ setting was crucial in ([36], [22]). And the higher rank Morse Lemma was heavily used in [32] to show that the visual metric defined in terms of the higher rank Gromov product has the desired properties.

On the other hand, the coarse feature of a Gromov hyperbolic space does not make visual metrics as good as in the $\text{CAT}(-1)$ setting, and the higher rank Morse lemma is not available in the generality of our setting. To overcome this difficulty, we conduct a detailed investigation of the coarse geometry in Gromov hyperbolic spaces to make the visual metric defined on the Gromov boundary work in higher rank settings, together with a simultaneous sharp-control on the other Busemann function on the higher rank symmetric space (see Remark 7.3).

Finally, we deduce the ergodicity of the horospherical foliation for hyper-transverse subgroups with respect to Burger-Roblin measures based on our investigation on the essential subgroup.

Organization. In Section 2, we review basic notions and properties of semisimple real algebraic groups and their boundaries. In Section 3 we introduce the notion of essential subgroups for conformal measures and discuss how the essential subgroup of a given conformal measure detects its singularity among conformal measures. In Section 4, we define self-joining subgroups and graph-conformal measures, and describe their key role in studying rigidity questions. We introduce the notion of Myrberg limit set in higher rank in Section 5. We prove that the Myrberg limit set of a self-joining has full mass with respect to the graph-conformal measure. Section 6 is devoted to the discussion on quantitative aspects of the geometry of Gromov hyperbolic spaces. In Section 7, we show that the essential subgroup for a graph-conformal measure is the whole \mathfrak{a}_θ under the Zariski dense hypothesis on the self-joining. In Section 8, we establish the singularity of graph-conformal measures among conformal measures, and provide the proof of Theorem 1.4. In Section 9, we prove more stronger rigidity statements (Theorem 1.8, Theorem 1.9) for deformations of transverse representations. The ergodicity of horospherical foliations is proved in Section 10.

Acknowledgements. I would like to thank my advisor Professor Hee Oh for her encouragement and many helpful conversations.

2. PRELIMINARIES

Let G be a connected semisimple real algebraic group. We use the notations and terminology introduced in the introduction. We denote by $\mathcal{W} = N_K(A)/C_K(A)$ the Weyl group, where $N_K(A)$ and $C_K(A)$ are the normalizer and centralizer of A in K respectively. Fixing a left G -invariant and right K -invariant Riemannian metric on G induces a \mathcal{W} -invariant norm

on \mathfrak{a} , which we denote by $\|\cdot\|$. We also denote by d the induced left G -invariant metric on the symmetric space $X := G/K$ and by $o \in X$ the point corresponding to the coset $[K]$.

Recall that we choose a closed positive Weyl chamber \mathfrak{a}^+ of \mathfrak{a} and set $A^+ = \exp \mathfrak{a}^+$. The Cartan projection $\mu : G \rightarrow \mathfrak{a}^+$ is defined to be such that $g \in K \exp(\mu(g))K$ for all $g \in G$.

Lemma 2.1. [2, Lemma 4.6] *For any compact subset $Q \subset G$, there exists $C = C(Q) > 0$ such that for all $g \in G$,*

$$\sup_{q_1, q_2 \in Q} \|\mu(q_1 g q_2) - \mu(g)\| \leq C.$$

Let Φ^+ be the set of all positive roots and $\Pi \subset \Phi^+$ the set of all simple roots for $(\mathfrak{g}, \mathfrak{a}^+)$. We fix an element

$$w_0 \in N_K(A)$$

representing the longest Weyl element. This induces an involution

$$i := -\text{Ad}_{w_0} : \mathfrak{a} \rightarrow \mathfrak{a}$$

preserving \mathfrak{a}^+ , called the opposition involution. This also induces a map $\Phi \rightarrow \Phi$ preserving Π , for which we use the same notation i . We have

$$(2.1) \quad \mu(g^{-1}) = i(\mu(g)) \quad \text{for all } g \in G.$$

In the rest of the section, fix a non-empty subset $\theta \subset \Pi$. Let P_θ denote a standard parabolic subgroup of G corresponding to θ ; that is, P_θ is generated by MA and all root subgroups U_α , $\alpha \in \Phi^+ \cup [\Pi - \theta]$ where $[\Pi - \theta]$ denotes the set of all roots in Φ which are \mathbb{Z} -linear combinations of $\Pi - \theta$. Hence $P_\Pi = P$. The subgroup P_θ is equal to its own normalizer; for $g \in G$, $gP_\theta g^{-1} = P_\theta$ if and only if $g \in P_\theta$. Let

$$\begin{aligned} \mathfrak{a}_\theta &= \bigcap_{\alpha \in \Pi - \theta} \ker \alpha, & \mathfrak{a}_\theta^+ &= \mathfrak{a}_\theta \cap \mathfrak{a}^+, \\ A_\theta &= \exp \mathfrak{a}_\theta, & \text{and } A_\theta^+ &= \exp \mathfrak{a}_\theta^+. \end{aligned}$$

Let

$$p_\theta : \mathfrak{a} \rightarrow \mathfrak{a}_\theta$$

denote the projection invariant under $w \in \mathcal{W}$ fixing \mathfrak{a}_θ pointwise. We also write $\mu_\theta := p_\theta \circ \mu$. We denote by $\mathfrak{a}_\theta^* = \text{Hom}(\mathfrak{a}_\theta, \mathbb{R})$ the dual space of \mathfrak{a}_θ . It can be identified with the subspace of \mathfrak{a}^* which is p_θ -invariant: $\mathfrak{a}_\theta^* = \{\psi \in \mathfrak{a}^* : \psi \circ p_\theta = \psi\}$; so for $\theta \subset \theta'$, we have $\mathfrak{a}_\theta^* \subset \mathfrak{a}_{\theta'}^*$.

Let L_θ be the centralizer of A_θ ; it is a Levi subgroup of P_θ and $P_\theta = L_\theta N_\theta$ where $N_\theta = R_u(P_\theta)$ is the unipotent radical of P_θ . We set $M_\theta = K \cap P_\theta \subset L_\theta$. We may then write $L_\theta = A_\theta S_\theta$ where S_θ is an almost direct product of a connected semisimple real algebraic subgroup and a compact subgroup. We omit the subscript when $\theta = \Pi$.

The θ -boundary \mathcal{F}_θ . The Furstenberg boundary is defined as the quotient $\mathcal{F} = G/P = G/P_\Pi$. The θ -boundary is defined similarly:

$$\mathcal{F}_\theta = G/P_\theta.$$

Let

$$\pi_\theta : \mathcal{F} \rightarrow \mathcal{F}_\theta$$

denote the canonical projection map given by $gP \mapsto gP_\theta$, $g \in G$. We set

$$\xi_\theta = [P_\theta] \in \mathcal{F}_\theta.$$

By the Iwasawa decomposition $G = KP = KAN$, the subgroup K acts transitively on \mathcal{F}_θ , and hence

$$\mathcal{F}_\theta \simeq K/M_\theta.$$

Points in general position. Let P_θ^+ be the standard parabolic subgroup of G opposite to P_θ such that $P_\theta \cap P_\theta^+ = L_\theta$. We have $P_\theta^+ = w_0 P_{i(\theta)} w_0^{-1}$ and hence

$$\mathcal{F}_{i(\theta)} = G/P_\theta^+.$$

In particular, if θ is symmetric in the sense that $\theta = i(\theta)$, then $\mathcal{F}_\theta = G/P_\theta^+$. The G -orbit of (P_θ, P_θ^+) is the unique open G -orbit in $G/P_\theta \times G/P_\theta^+$ under the diagonal G -action.

Definition 2.2. Two elements $\xi \in \mathcal{F}_\theta$ and $\eta \in \mathcal{F}_{i(\theta)}$ are said to be in general position if $(\xi, \eta) \in G \cdot (P_\theta, w_0 P_{i(\theta)}) = G \cdot (P_\theta, P_\theta^+)$, i.e., $\xi = gP_\theta$ and $\eta = gw_0 P_{i(\theta)}$ for some $g \in G$.

We set

$$\mathcal{F}_\theta^{(2)} = \{(\xi, \eta) \in \mathcal{F}_\theta \times \mathcal{F}_{i(\theta)} : \xi, \eta \text{ are in general position}\},$$

which is the unique open G -orbit in $\mathcal{F}_\theta \times \mathcal{F}_{i(\theta)}$.

Jordan projection. An element $g \in G$ is loxodromic if

$$g = hamh^{-1}$$

for some $a \in \text{int } A^+$, $m \in M$ and $h \in G$. The Jordan projection of g is defined to be

$$\lambda(g) := \log a \in \text{int } \mathfrak{a}^+.$$

The attracting fixed point of g in \mathcal{F} is given by

$$y_g := hP \in \mathcal{F};$$

for any $\xi \in \mathcal{F}$ in general position with $y_{g^{-1}}$, the sequence $g^\ell \xi$ converges to y_g as $\ell \rightarrow \infty$. We also set

$$\lambda_\theta(g) := p_\theta(\lambda(g)) \in \text{int } \mathfrak{a}_\theta^+ \quad \text{and} \quad y_g^\theta := \pi_\theta(y_g) \in \mathcal{F}_\theta.$$

Since $g^{-1} = (hw_0)(w_0^{-1}a^{-1}w_0)(w_0^{-1}m^{-1}w_0)(hw_0)^{-1}$ and $w_0^{-1}aw_0 \in \text{int } \mathfrak{a}^+$ and $w_0^{-1}mw_0 \in M$ as well, it follows that y_g^θ and $y_{g^{-1}}^{i(\theta)}$ are in general position.

Let $\Delta < G$ be a discrete subgroup. We write $\lambda(\Delta)$ for the set of all Jordan projections of loxodromic elements of Δ . The following result is due to Benoist [3].

Theorem 2.3. *If $\Delta < G$ is Zariski dense, then $\lambda(\Delta)$ generates a dense subgroup of \mathfrak{a} . In particular, $\lambda_\theta(\Delta)$ generates a dense subgroup of \mathfrak{a}_θ .*

Convergence in $G \cup \mathcal{F}_\theta$. We consider the following notion of convergence of a sequence in G to an element of \mathcal{F}_θ . We say that for a sequence $g_i \in G$, $g_i \rightarrow \infty$ (or $g_i o \rightarrow \infty$) θ -regularly if $\min_{\alpha \in \theta} \alpha(\mu(g_i)) \rightarrow \infty$ as $i \rightarrow \infty$.

Definition 2.4. For a sequence $g_i \in G$ and $\xi \in \mathcal{F}_\theta$, we write $\lim_{i \rightarrow \infty} g_i = \lim_{i \rightarrow \infty} g_i o = \xi$ and say g_i (or $g_i o \in X$) converges to ξ if

- $g_i \rightarrow \infty$ θ -regularly; and
- $\lim_{i \rightarrow \infty} \kappa_{g_i} P_\theta = \xi$ in \mathcal{F}_θ for some $\kappa_{g_i} \in K$ such that $g_i \in \kappa_{g_i} A^+ K$.

We recall the lemma that we will use in later sections.

Lemma 2.5. [26, Lemma 2.4] *Consider a sequence $g_i = k_i a_i h_i^{-1}$ where $k_i \in K$, $a_i \in A^+$, and $h_i \in G$. Suppose that $k_i \rightarrow k_0 \in K$, $h_i \rightarrow h_0 \in G$, and $\min_{\alpha \in \theta} \alpha(\log a_i) \rightarrow \infty$, as $i \rightarrow \infty$. Then for any ξ in general position with $h_0 P_\theta^+$, we have*

$$\lim_{i \rightarrow \infty} g_i \xi = k_0 \xi_\theta.$$

Let $p, q \in X$ and $R > 0$. The shadow of the ball

$$B(q, R) = \{z \in X : d(z, q) < R\}$$

viewed from p is defined as follows:

$$O_R^\theta(p, q) := \{g P_\theta \in \mathcal{F}_\theta : g A^+ o \cap B(q, r) \neq \emptyset\}$$

where $g \in G$ satisfies $p = go$. The shadow of $B(q, R)$ viewed from $\eta \in \mathcal{F}_{i(\theta)}$ can also be defined:

$$O_R^\theta(\eta, q) := \{h P_\theta \in \mathcal{F}_\theta : h w_0 P_{i(\theta)} = \eta, h o \in B(q, r)\}.$$

We say that a sequence $g_i \in G$ (or $g_i o \in X$) converges to $\xi \in \mathcal{F}_\theta$ conically if $g_i \rightarrow \xi$ in the sense of Definition 2.4 and there exists $R > 0$ such that $\xi \in O_R^\theta(o, g_i o)$ for all $i \geq 1$.

Lemma 2.6 ([21, Lemma 5.35] (see also [26, Lemma 9.8])). *Let $g_i \in G$ be a sequence such that $g_i \rightarrow \xi \in \mathcal{F}_\theta$. Then the following are equivalent:*

- (1) *The convergence $g_i \rightarrow \xi$ is conical.*
- (2) *For any $\eta \in \mathcal{F}_{i(\theta)}$ such that $(\xi, \eta) \in \mathcal{F}_\theta^{(2)}$, the sequence $g_i^{-1}(\xi, \eta)$ is precompact in $\mathcal{F}_\theta^{(2)}$.*
- (3) *For some $\eta \in \mathcal{F}_{i(\theta)}$ such that $(\xi, \eta) \in \mathcal{F}_\theta^{(2)}$, the sequence $g_i^{-1}(\xi, \eta)$ is precompact in $\mathcal{F}_\theta^{(2)}$.*

The shadows vary continuously:

Lemma 2.7. [25, Lemma 3.3] *Let $p \in X$, $q_i \in X$ a sequence converging to $\eta \in \mathcal{F}_{i(\theta)}$ and $r > 0$. Then for any $p \in X$ and $0 < \varepsilon < r$, we have for all large enough i that*

$$O_{r-\varepsilon}^\theta(q_i, p) \subset O_r^\theta(\eta, p) \subset O_{r+\varepsilon}^\theta(q_i, p).$$

Limit set. For a discrete subgroup $\Delta < G$, its limit set is defined as follows:

Definition 2.8 (Limit set). The limit set $\Lambda_\Delta^\theta \subset \mathcal{F}_\theta$ is defined as the set of all accumulation points of $\Delta(o)$ in \mathcal{F}_θ , that is,

$$\Lambda_\Delta^\theta = \left\{ \lim_{i \rightarrow \infty} g_i o \in \mathcal{F}_\theta : g_i \in \Delta \right\}.$$

This may be an empty set for a general discrete subgroup. However, we have the following result of Benoist for Zariski dense subgroups ([2, Section 3.6], [32, Lemma 2.15]):

Theorem 2.9. *If $\Delta < G$ is Zariski dense, then Λ_Δ^θ is the unique Δ -minimal subset of \mathcal{F}_θ and the set of all attracting fixed points of loxodromic elements of Δ is dense in Λ_Δ^θ .*

3. BUSEMANN MAPS, CONFORMAL MEASURES AND ESSENTIAL SUBGROUPS

Let G be a connected semisimple real algebraic group and fix a non-empty subset $\theta \subset \Pi$. We continue notations from Section 2. In this section, we introduce conformal measures and essential subgroups, and see how they are related.

Busemann maps. For $g \in G$ and $\xi = [k] \in K/M = \mathcal{F}$, the Iwasawa cocycle $H(g, \xi)$ is defined as the \mathfrak{a} -component of the Iwasawa decomposition of gk so that

$$gk \in K \exp(H(g, \xi))N.$$

The higher rank Busemann maps are defined as follows:

Definition 3.1 (Busemann map). The \mathfrak{a} -valued Busemann map $\beta : \mathcal{F} \times G \times G \rightarrow \mathfrak{a}$ is now defined as follows: for $\xi \in \mathcal{F}$ and $g, h \in G$,

$$\beta_\xi(g, h) := H(g^{-1}, \xi) - H(h^{-1}, \xi).$$

The \mathfrak{a}_θ -valued Busemann map $\beta^\theta : \mathcal{F}_\theta \times G \times G \rightarrow \mathfrak{a}_\theta$ is defined as follows: for $\xi \in \mathcal{F}_\theta$ and $g, h \in G$,

$$\beta_\xi^\theta(g, h) := p_\theta(\beta_{\tilde{\xi}}(g, h))$$

where $\tilde{\xi} \in \pi_\theta^{-1}(\xi) \in \mathcal{F}$. This is well-defined [35, Section 6].

Observe that the Busemann map is continuous in all three variables. For $\xi \in \mathcal{F}$, $g \in G$ and $k \in K$, we have $H((gk)^{-1}, \xi) = H(g^{-1}, \xi)$. Hence we can also define the Busemann map $\beta^\theta : \mathcal{F}_\theta \times X \times X \rightarrow \mathfrak{a}$ as

$$\beta_\xi^\theta(go, ho) := \beta_\xi^\theta(g, h) \quad \text{for } \xi \in \mathcal{F}_\theta, g, h \in G.$$

The following was proved in [32] for $\theta = \Pi$. Since the \mathfrak{a}_θ -valued Busemann map is defined as the p_θ -image of the \mathfrak{a} -valued Busemann map, the same is true for general θ :

Lemma 3.2. [32, Lemma 3.5] *For a loxodromic $g \in G$, we have*

$$\beta_{y_g^\theta}^\theta(p, gp) = \lambda_\theta(g) \quad \text{for all } p \in X.$$

Busemann maps are compatible to Cartan projections in shadows:

Lemma 3.3. [32, Lemma 5.7] *There exists $\kappa > 0$ such that for any $g, h \in G$ and $r > 0$, we have*

$$\sup_{\xi \in O_r^\theta(go, ho)} \|\beta_\xi^\theta(go, ho) - \mu_\theta(g^{-1}h)\| \leq \kappa r.$$

Corollary 3.4. *Let $g_i \in G$ be a sequence such that $g_i o \rightarrow \eta \in \mathcal{F}_i(\theta)$. For any $p \in X$ and $r, \varepsilon > 0$, we have*

$$\sup_{\xi, \xi' \in O_r^\theta(\eta, p)} \|\beta_\xi^\theta(g_i o, p) - \beta_{\xi'}^\theta(g_i o, p)\| \leq 2\kappa(r + \varepsilon) \quad \text{for all large } i$$

where κ is given by Lemma 3.3.

Proof. By Lemma 2.7, we have $O_r^\theta(\eta, p) \subset O_{r+\varepsilon}^\theta(g_i o, p)$ for all large i . Letting $h \in G$ be such that $p = ho$, we have that for any $\xi' \in O_r^\theta(\eta, p)$, both $\|\beta_{\xi'}^\theta(g_i o, p) - \mu_\theta(g_i^{-1}h)\|$ and $\|\beta_{\xi'}^\theta(g_i o, p) - \mu_\theta(g_i^{-1}h)\|$ are bounded by $\kappa(r + \varepsilon)$ by Lemma 3.3. Now the claim follows from the triangle inequality. \square

Conformal measures. Let $\Delta < G$ be a discrete subgroup. The notion of higher rank conformal measures for Δ is defined in terms of \mathfrak{a}_θ -valued Busemann maps and linear forms on \mathfrak{a}_θ .

Definition 3.5 (Conformal measures). A Borel probability measure ν_o on \mathcal{F}_θ is called a Δ -conformal measure (with respect to o) if there exists a linear form $\psi \in \mathfrak{a}_\theta^*$ such that for all $\eta \in \mathcal{F}_\theta$ and $g \in \Delta$,

$$\frac{dg_* \nu_o}{d\nu_o}(\eta) = e^{\psi(\beta_\eta^\theta(o, go))}.$$

In this case, we say ν_o is a (Δ, ψ) -conformal measure. For $p \in X$, $d\nu_p(\eta) = e^{\psi(\beta_\eta^\theta(o, p))} d\nu_o(\eta)$ is a (Δ, ψ) -conformal measure with respect to p .

The set of values $\{\beta_\eta^\theta(o, go) \in \mathfrak{a}_\theta : g \in \Delta, \eta \in \text{supp } \nu_o\}$ may not be large enough to distinguish Δ -conformal measure classes by determining the linear form to which ν_o is associated; in general, there may be multiple linear forms associated to the same conformal measure class.

Definition 3.6 (Divergence type). We say that a (Δ, ψ) -conformal measure ν is of *divergence type* if $\psi \in \mathfrak{a}_\theta^*$ is (Γ, θ) -proper and $\sum_{g \in \Delta} e^{-\psi(\mu_\theta(g))} = \infty$.

Essential subgroups and Singularity of conformal measures. The notion of essential subgroups was introduced by Schmidt [38] (see also [36]) in order to study the ergodic properties of horospherical actions. Its higher rank analogue, when $\theta = \Pi$, was studied in [32] to show the ergodicity of horospherical foliations for Anosov subgroups with respect to a minimal parabolic subgroup. We consider essential subgroups for general θ , and also use them as tools to detect the singularity between two conformal measures.

Definition 3.7 (Essential subgroup for ν). For a Δ -conformal measure ν on \mathcal{F}_θ with respect to o , we define the subset $\mathbf{E}_\nu^\theta(\Delta) \subset \mathfrak{a}_\theta$ as follows: $u \in \mathbf{E}_\nu^\theta(\Delta)$ if for any Borel subset $B \subset \mathcal{F}_\theta$ with $\nu(B) > 0$ and any $\varepsilon > 0$, there exists $g \in \Delta$ such that

$$\nu(B \cap gB \cap \{\xi \in \mathcal{F}_\theta : \|\beta_\xi^\theta(o, go) - u\| < \varepsilon\}) > 0.$$

It is easy to see that $\mathbf{E}_\nu^\theta(\Delta)$ is a closed subgroup of \mathfrak{a}_θ . We call $\mathbf{E}_\nu^\theta(\Delta)$ the essential subgroup for ν .

The following proposition is one of key ingredients of this paper. Although it was proved in [32, Lemma 10.21] (see also [22, Proposition 3.6]) for $\theta = \Pi$, the same proof works for general θ :

Proposition 3.8. *For $i = 1, 2$, let ν_i be a (Δ, ψ_i) -conformal measure on \mathcal{F}_θ for some $\psi_i \in \mathfrak{a}_\theta^*$. If $\nu_2 \ll \nu_1$, then*

$$\psi_1(w) = \psi_2(w) \quad \text{for all } w \in \mathbf{E}_{\nu_1}^\theta(\Delta).$$

In particular, if $\mathbf{E}_{\nu_1}^\theta(\Delta) = \mathfrak{a}_\theta$, then $\nu_2 \ll \nu_1$ implies $\psi_1 = \psi_2$.

4. GRAPH-CONFORMAL MEASURES FOR SELF-JOININGS

In this section, we review the notion of self-joinings and graph-conformal measures, introduced by Kim-Oh ([23], [24], [22]), which play key roles in studying rigidity problems. For $i = 1, 2$, let G_i be a connected semisimple real algebraic group with the associated Riemannian symmetric space (X_i, d_i) , and write $\mathfrak{g}_i := \text{Lie } G_i$. Let (X, d) be the Riemannian product $(X_1 \times X_2, \sqrt{d_1^2 + d_2^2})$. Set

$$G = G_1 \times G_2$$

so that its Lie algebra is $\mathfrak{g} := \mathfrak{g}_1 \oplus \mathfrak{g}_2$. Then G acts by isometries on X . For $i = 1, 2$, we use the same notations for G_i as we did for G but with a subscript i . For $\square \in \{A, M, N, P, K, o\}$, we consider the corresponding objects for G by setting

$$\square = \square_1 \times \square_2.$$

In particular, $A = A_1 \times A_2$. Let $A^+ = A_1^+ \times A_2^+$. Let \mathfrak{a} denote the Lie algebra of A , and $\mathfrak{a}^+ = \log A^+$. We note that

$$\mathfrak{a} = \mathfrak{a}_1 \oplus \mathfrak{a}_2 \quad \text{and} \quad \mathfrak{a}^+ = \mathfrak{a}_1^+ \oplus \mathfrak{a}_2^+,$$

where $\mathfrak{a}_i = \text{Lie } A_i$ and $\mathfrak{a}_i^+ = \text{Lie } A_i^+$ for $i = 1, 2$.

For each $i = 1, 2$, let Π_i be the set of all simple roots for $(\mathfrak{g}_i, \mathfrak{a}_i^+)$ and fix a non-empty subset $\theta_i \subset \Pi_i$ in the rest of the paper. We set $\Pi := \Pi_1 \cup \Pi_2$ which is the set of all simple roots for $(\mathfrak{g}, \mathfrak{a}^+)$ and

$$\theta := \theta_1 \cup \theta_2.$$

Then we have

$$\mathfrak{a}_\theta = \mathfrak{a}_{\theta_1} \oplus \mathfrak{a}_{\theta_2}, \quad P_\theta = P_{\theta_1} \times P_{\theta_2} \quad \text{and} \quad \mathcal{F}_\theta = \mathcal{F}_{\theta_1} \times \mathcal{F}_{\theta_2}.$$

Let $\Gamma < G_1$ be a Zariski dense discrete subgroup with the limit set $\Lambda^{\theta_1} \subset \mathcal{F}_{\theta_1}$. Let $\rho : \Gamma \rightarrow G_2$ be a discrete faithful Zariski dense representation.

Definition 4.1 (Self-joining). We define the self-joining of Γ via ρ as

$$\Gamma_\rho := (\text{id} \times \rho)(\Gamma) = \{(g, \rho(g)) \in G : g \in \Gamma\},$$

which is a discrete subgroup of G .

One key feature of a self-joining subgroup is that the rigidity question on ρ can be translated to a Zariski density question on the self-joining.

Lemma 4.2. [15] *Suppose that G_1 and G_2 are simple. Then the self-joining $\Gamma_\rho < G$ is not Zariski dense if and only if ρ extends to a Lie group isomorphism $G_1 \rightarrow G_2$.*

Boundary map. In the rest of this section, we assume that there exists a ρ -equivariant continuous map

$$f : \Lambda^{\theta_1} \rightarrow \mathcal{F}_{\theta_2}.$$

We will not assume that f is injective, mentioned otherwise. When it is injective, we call it a ρ -boundary map.

A ρ -boundary map is unique when it exists. First observe that, since Λ^{θ_1} (resp. $\Lambda_{\rho(\Gamma)}^{\theta_2}$) is the unique Γ (resp. $\rho(\Gamma)$) minimal subset of \mathcal{F}_{θ_1} (resp. \mathcal{F}_{θ_2}) (Theorem 2.9), it follows from the equivariance of f that

$$f(\Lambda^{\theta_1}) = \Lambda_{\rho(\Gamma)}^{\theta_2}.$$

The uniqueness of a boundary map was proved in [22, Lemma 4.5] for $\theta = \Pi$ and the same proof works for general θ .

Lemma 4.3 (Uniqueness). *If $g \in \Gamma$ and $\rho(g)$ are both loxodromic, then*

$$f(y_g^{\theta_1}) = y_{\rho(g)}^{\theta_2}.$$

In particular, when G_1 and G_2 are simple, f is the unique ρ -equivariant continuous map $\Lambda^{\theta_1} \rightarrow \mathcal{F}_{\theta_2}$.

In terms of the boundary map, the limit set of the self-joining Γ_ρ in \mathcal{F}_θ is as follows:

$$\Lambda_\rho^\theta := (\text{id} \times f)(\Lambda^{\theta_1}) = \{(\xi, f(\xi)) \in \mathcal{F}_\theta : \xi \in \Lambda^{\theta_1}\}.$$

Graph-conformal measures. Recall that for each $i = 1, 2$,

$$p_{\theta_i} : \mathfrak{a} \rightarrow \mathfrak{a}_{\theta_i}$$

is the projection invariant under all Weyl elements fixing \mathfrak{a}_{θ_i} pointwise. Restricting on \mathfrak{a}_θ , we may regard p_{θ_i} as the projection from \mathfrak{a}_θ as well. Abusing notations, we also denote the restriction $\mathfrak{a}_\theta \rightarrow \mathfrak{a}_{\theta_i}$ by p_{θ_i} for $i = 1, 2$. For a linear form ψ_i on $\mathfrak{a}_{\theta_i}^*$, we define a linear form $\sigma_{\psi_i} \in \mathfrak{a}_\theta^*$ by

$$\sigma_{\psi_i} := \psi_i \circ p_{\theta_i};$$

in other words, $\sigma_{\psi_i}(u_1, u_2) = \psi_i(u_i)$ for all $(u_1, u_2) \in \mathfrak{a}_{\theta_1} \oplus \mathfrak{a}_{\theta_2}$. Recall that $o_i = [K_i] \in X_i$ for $i = 1, 2$ and $o = (o_1, o_2) \in X$.

The following is a key observation on the relation between Γ -conformal measures and Γ_ρ -conformal measures. It was proved in [22, Proposition 4.6, Corollary 4.7, Lemma 4.10] for $\theta = \Pi$; the same proof works for general θ :

Proposition 4.4. *Let $\psi_i \in \mathfrak{a}_{\theta_i}^*$ for $i = 1, 2$.*

- (1) *If ν_{ψ_1} is a (Γ, ψ_1) -conformal measure on Λ^{θ_1} with respect to o_1 , then $(\text{id} \times f)_* \nu_{\psi_1}$ is a $(\Gamma_\rho, \sigma_{\psi_1})$ -conformal measure on Λ_ρ^θ with respect to o .*
- (2) *Any $(\Gamma_\rho, \sigma_{\psi_1})$ -conformal measure on Λ_ρ^θ with respect to o is of the form $(\text{id} \times f)_* \nu_{\psi_1}$ for some (Γ, ψ_1) -conformal measure ν_{ψ_1} on Λ^{θ_1} .*
- (3) *If ν_{ψ_1} is the unique (Γ, ψ_1) -conformal measure on Λ^{θ_1} , then $(\text{id} \times f)_* \nu_{\psi_1}$ is the unique $(\Gamma_\rho, \sigma_{\psi_1})$ -conformal measure on Λ_ρ^θ with respect to o ; in particular, $(\text{id} \times f)_* \nu_{\psi_1}$ is Γ_ρ -ergodic.*
- (4) *Let ν_{ψ_1} and ν_{ψ_2} be Γ -conformal and $\rho(\Gamma)$ -conformal measures on Λ^{θ_1} and $\Lambda_{\rho(\Gamma)}^{\theta_2}$ respectively. If f is injective, then $(f^{-1} \times \text{id})_* \nu_{\psi_2}$ is a $(\Gamma_\rho, \sigma_{\psi_2})$ -conformal measure, and we have*

$$(f^{-1} \times \text{id})_* \nu_{\psi_2} \ll (\text{id} \times f)_* \nu_{\psi_1} \text{ if and only if } \nu_{\psi_2} \ll f_* \nu_{\psi_1}.$$

The notion of graph-conformal measure was first introduced in our earlier work with Oh [22]:

Definition 4.5 (Graph-conformal measures). By a graph-conformal measure of Γ_ρ , we mean a (conformal) measure of the form

$$\nu_\rho := (\text{id} \times f)_* \nu$$

for some Γ -conformal measure ν on Λ^{θ_1} .

Note that we used the notation ν_{graph} for the graph-conformal measure in [22]. Using this terminology, Proposition 4.4(1)-(2) can be reformulated as follows:

Proposition 4.6. *Let $\sigma \in \mathfrak{a}_\theta^*$ be a linear form which factors through \mathfrak{a}_{θ_1} . A (Γ_ρ, σ) -conformal measure on Λ_ρ^θ is a graph-conformal measure of Γ_ρ , and conversely, any graph-conformal measure of Γ_ρ is of such a form.*

5. MYRBERG LIMIT SETS

Let G be a connected semisimple real algebraic group and $X = G/K$ the associated symmetric space. Recall the choice of the basepoint $o = [K] \in X$. For a discrete subgroup $\Delta < G$, the Myrberg limit set is defined as follows:

Definition 5.1 (Myrberg limit set). We say that $\xi \in \Lambda_\Delta^\theta$ is a Myrberg limit point of Δ if for any $\xi_0 \in \Lambda_\Delta^\theta$ and $\eta_0 \in \Lambda_\Delta^{i(\theta)}$ in general position, there exists a sequence $g_i \in \Delta$ such that $g_i \xi \rightarrow \xi_0$ and $g_i o \rightarrow \eta_0$ as $i \rightarrow \infty$.

We call the set of all Myrberg limit points of Δ the *Myrberg limit set* of Δ and denote by $\Lambda_{\Delta, M}^\theta \subset \mathcal{F}_\theta$.

In this section, we show that the Myrberg limit sets of transverse subgroups and their self-joinings have full measures with respect to conformal measures of divergence type and their associated graph-conformal measures.

Definition 5.2. A discrete subgroup $\Delta < G$ is called θ -transverse if

- it is θ -regular, i.e., $\liminf_{g \in \Delta} \alpha(\mu(g)) = \infty$ for all $\alpha \in \theta$; and
- it is θ -antipodal, i.e., any two distinct $\xi, \eta \in \Lambda^{\theta \cup i(\theta)}$ are in general position.

Since $\mu(g^{-1}) = i(\mu(g))$ for all $g \in G$, the θ -regularity is equivalent to the $i(\theta)$ -regularity, and hence Δ is θ -regular if and only if it is $\theta \cup i(\theta)$ -regular. Moreover, by [26, Lemma 9.5], the θ -antipodality implies that the canonical projections $\Lambda_\Delta^{\theta \cup i(\theta)} \rightarrow \Lambda_\Delta^\theta$ and $\Lambda_\Delta^{\theta \cup i(\theta)} \rightarrow \Lambda_\Delta^{i(\theta)}$ are Δ -equivariant homeomorphisms.

Myrberg limit sets of transverse subgroups and their self-joinings.

For $i = 1, 2$, let G_i be a connected semisimple real algebraic group and X_i the associated symmetric space and fix a non-empty subset $\theta_i \subset \Pi_i$. Let

$$\Gamma < G_1$$

be a Zariski dense θ_1 -transverse subgroup with limit set

$$\Lambda^{\theta_1} := \Lambda_\Gamma^{\theta_1} \subset \mathcal{F}_{\theta_1}.$$

Let

$$\rho : \Gamma \rightarrow G_2$$

be a Zariski dense θ_2 -regular faithful representation with a pair of ρ -equivariant continuous maps $f : \Lambda^{\theta_1} \rightarrow \Lambda_{\rho(\Gamma)}^{\theta_2}$ and $f_i : \Lambda^{i(\theta_1)} \rightarrow \Lambda_{\rho(\Gamma)}^{i(\theta_2)}$.

Set

$$G = G_1 \times G_2, \quad X = X_1 \times X_2, \quad \text{and} \quad \theta = \theta_1 \cup \theta_2.$$

We keep the notations introduced in Section 4. We denote by

$$\Lambda_\rho^\theta \subset \mathcal{F}_\theta$$

the limit set of the self-joining Γ_ρ . We simply write

$$\Lambda_M^{\theta_1} := \Lambda_{\Gamma, M}^{\theta_1} \subset \mathcal{F}_{\theta_1} \quad \text{and} \quad \Lambda_{\rho, M}^\theta := \Lambda_{\Gamma_\rho, M}^\theta \subset \mathcal{F}_\theta$$

for the Myrberg limit sets of Γ and Γ_ρ respectively. The main goal of this section is to prove the following:

Theorem 5.3. *Let ν be a Γ -conformal measure on \mathcal{F}_{θ_1} of divergence type, and $\nu_\rho = (\text{id} \times f)_* \nu$ the associated graph-conformal measure for Γ_ρ on \mathcal{F}_θ . Then*

$$\nu(\Lambda_M^{\theta_1}) = 1 \quad \text{and} \quad \nu_\rho(\Lambda_{\rho, M}^\theta) = 1.$$

Remark 5.4. The claim $\nu(\Lambda_M^{\theta_1}) = 1$ was proved by Tukia [42, Theorem 4A] when $\text{rank } G_1 = 1$, and by Lee-Oh [32] when Γ is Π -Anosov.

Ergodic properties of divergence-type conformal measures. To prove Theorem 5.3, we use the ergodic properties of conformal measures of divergence type. Recall that a linear form $\psi \in \mathfrak{a}_{\theta_1}^*$ is called (Γ, θ_1) -proper if the map $\psi \circ \mu_{\theta_1}|_\Gamma : \Gamma \rightarrow [-\varepsilon, \infty)$ is proper for some $\varepsilon > 0$. For a (Γ, θ_1) -proper $\psi \in \mathfrak{a}_{\theta_1}^*$, the abscissa of convergence $\delta_\psi \in (0, \infty]$ of the ψ -Poincaré series $\sum_{g \in \Gamma} e^{-s\psi(\mu_{\theta_1}(g))}$ is well-defined [26, Lemma 4.2]. Moreover, if there exists a (Γ, ψ) -conformal measure, then $\delta_\psi \leq 1$ [26, Theorem 1.5]. Whether or not the ψ -Poincaré series converges or diverges at $s = 1$ plays a crucial role:

Theorem 5.5. [26, Theorem 1.5] *Let $\psi \in \mathfrak{a}_{\theta_1}^*$ be a (Γ, θ_1) -proper form. If $\delta_\psi \leq 1$ and $\sum_{g \in \Gamma} e^{-\psi(\mu_{\theta_1}(g))} = \infty$, then there exists a unique (Γ, ψ) -conformal measure on \mathcal{F}_{θ_1} . Moreover, the unique (Γ, ψ) -conformal measure has the support Λ^{θ_1} . In particular, any Γ -conformal measure of divergence type is Γ -ergodic.*

Under the additional assumption on the support, the above statement was also proved in [13].

Recall that $\mathcal{F}_{\theta_1}^{(2)} \subset \mathcal{F}_{\theta_1} \times \mathcal{F}_{i(\theta_1)}$ is the set of all pairs in general position. Fixing a (Γ, θ_1) -proper form $\varphi \in \mathfrak{a}_{\theta_1}^*$, we consider the G_1 -action

$$(5.1) \quad g \cdot (\xi, \eta, s) = (g\xi, g\eta, s + \varphi(\beta_\xi^{\theta_1}(g^{-1}, e)))$$

for all $g \in G_1$ and $(\xi, \eta, s) \in \mathcal{F}_{\theta_1}^{(2)} \times \mathbb{R}$. Setting

$$\Lambda^{(2)} := (\Lambda^{\theta_1} \times \Lambda^{i(\theta_1)}) \cap \mathcal{F}_{\theta_1}^{(2)},$$

the subspace $\Lambda^{(2)} \times \mathbb{R} \subset \mathcal{F}_{\theta_1}^{(2)} \times \mathbb{R}$ is invariant under the Γ -action.

Theorem 5.6. [26, Theorem 9.2] *The action Γ on $\Lambda^{(2)} \times \mathbb{R}$ given by (5.1) is properly discontinuous and hence*

$$\Omega_\varphi := \Gamma \backslash \Lambda^{(2)} \times \mathbb{R}$$

is a locally compact second-countable Hausdorff space.

Let $\psi \in \mathfrak{a}_{\theta_1}^*$. For a pair of a (Γ, ψ) -conformal measure ν on Λ^{θ_1} and a $(\Gamma, \psi \circ i)$ -conformal measure ν_i on $\Lambda^{i(\theta_1)}$, we define a Radon measure $\tilde{m}_{\nu, \nu_i}^\varphi$ on $\Lambda^{(2)} \times \mathbb{R}$ as follows:

$$d\tilde{m}_{\nu, \nu_i}^\varphi(\xi, \eta, t) = e^{\psi(\beta_\xi^{\theta_1}(e, g) + i(\beta_\eta^{i(\theta_1)}(e, g)))} d\nu(\xi) d\nu_i(\eta) dt$$

where $g \in G_1$ is such that $(\xi, \eta) = (gP_{\theta_1}, gw_0P_{i(\theta_1)})$ and dt is the Lebesgue measure on \mathbb{R} . This is well-defined [26, Lemma 9.13]. The measure $d\tilde{m}_{\nu, \nu_1}^\varphi$ is left Γ -invariant and invariant under the translation on \mathbb{R} . Hence it descends to the \mathbb{R} -invariant Radon measure

$$\mathfrak{m}_{\nu, \nu_1}^\varphi$$

on Ω_φ , which we call the Bowen-Margulis-Sullivan measure associated to the pair (ν, ν_1) .

Let $\mathcal{M}_\psi^{\theta_1}$ (resp. $\mathcal{M}_{\psi \circ i}^{i(\theta_1)}$) be the set of all (Γ, ψ) -conformal measures on \mathcal{F}_{θ_1} (resp. $(\Gamma, \psi \circ i)$ -conformal measures on $\mathcal{F}_{i(\theta_1)}$). The following is the Hopf-Tsuji-Sullivan dichotomy for transverse subgroups: we also denote by $\Lambda_c^{\theta_1} \subset \mathcal{F}_{\theta_1}$ the conical set of Γ .

Theorem 5.7 ([26, Theorem 10.2], see also [13] and [39]). *Let $\Gamma < G_1$ be a Zariski dense θ_1 -transverse subgroup. The following are equivalent to each other.*

- (1) $\sum_{g \in \Gamma} e^{-\psi(\mu_{\theta_1}(g))} = \infty$ (resp. $\sum_{g \in \Gamma} e^{-\psi(\mu_{\theta_1}(g))} < \infty$);
- (2) $\nu(\Lambda_c^{\theta_1}) = 1$ (resp. $\nu(\Lambda_c^{\theta_1}) = 0$);
- (3) For any $(\nu, \nu_1) \in \mathcal{M}_\psi^{\theta_1} \times \mathcal{M}_{\psi \circ i}^{i(\theta_1)}$ and any (Γ, θ_1) -proper $\varphi \in \mathfrak{a}_{\theta_1}^*$, the \mathbb{R} -action on $(\Omega_\varphi, \mathfrak{m}_{\nu, \nu_1}^\varphi)$ is completely conservative and ergodic (resp. completely dissipative and non-ergodic).

Indeed, Theorem 5.3 adds one more item $\nu(\Lambda_M^{\theta_1}) = 1$ (resp. $\nu(\Lambda_M^{\theta_1}) = 0$) to the above dichotomy, since $\Lambda_M^{\theta_1} \subset \Lambda_c^{\theta_1}$ (Lemma 2.6).

Proof of Theorem 5.3. Note that we can regard ψ as a linear form on $\mathfrak{a}_{\theta_1 \cup i(\theta_1)}$ by precomposing with the projection $\mathfrak{a}_{\theta_1 \cup i(\theta_1)} \rightarrow \mathfrak{a}_{\theta_1}$, which is $(\Gamma, \theta_1 \cup i(\theta_1))$ -proper since $\psi(\mu_{\theta_1}(g)) = \psi(\mu_{\theta_1 \cup i(\theta_1)}(g))$ for all $g \in \Gamma$. By Theorem 5.7, ν is supported on the conical set of Γ , in particular, on Λ^{θ_1} . The canonical projection $\Lambda^{\theta_1 \cup i(\theta_1)} \rightarrow \Lambda^{\theta_1}$ is a Γ -equivariant homeomorphism [26, Lemma 9.5] and hence we can pull-back ν to $\Lambda^{\theta_1 \cup i(\theta_1)}$ so that ν can be considered as a (Γ, ψ) -conformal measure on $\Lambda^{\theta_1 \cup i(\theta_1)}$. Since θ_1 -transverse subgroups are $\theta_1 \cup i(\theta_1)$ -transverse, we may assume without loss of generality that $\theta_1 = i(\theta)$ by replacing θ_1 with $\theta_1 \cup i(\theta_1)$.

Myrberg limit set of Γ is ν -full. By [26, Theorem 1.5], it follows from the existence of ν that $\delta_\psi \leq 1$. Since $\sum_{g \in \Gamma} e^{-\psi(\mu_{\theta_1}(g))} = \infty$, we have $\delta_\psi = 1$. Hence ν is the unique (Γ, ψ) -conformal measure on \mathcal{F}_{θ_1} by Theorem 5.5, and is supported on Λ^{θ_1} as mentioned above. Moreover, since $\mu_{\theta_1}(g^{-1}) = i(\mu_{\theta_1}(g^{-1}))$, $\psi \circ i$ is also (Γ, θ_1) -proper, $\delta_{\psi \circ i} = 1$ and $\sum_{g \in \Gamma} e^{-(\psi \circ i)(\mu_{\theta_1}(g))} = \infty$. Hence by Theorem 5.5 and Theorem 5.7, there exists a unique $(\Gamma, \psi \circ i)$ -conformal measure ν_1 on \mathcal{F}_{θ_1} and is supported on Λ^{θ_1} as well.

Now we are able to consider the measure space $(\Omega_\varphi, \mathfrak{m}_{\nu, \nu_1}^\varphi)$ by fixing a (Γ, θ_1) -proper $\varphi \in \mathfrak{a}_{\theta_1}^*$. By Theorem 5.7, the \mathbb{R} -action on $(\Omega_\varphi, \mathfrak{m}_{\nu, \nu_1}^\varphi)$ is completely conservative and ergodic, and hence $\mathfrak{m}_{\nu, \nu_1}^\varphi$ -a.e. \mathbb{R}_+ -orbit is dense.

In other words, for $\nu \otimes \nu_1 \otimes dt$ -a.e. $(\xi, \eta, t) \in \Lambda^{(2)} \times \mathbb{R}$, its \mathbb{R}_+ -orbit is dense in Ω_φ . Fix one such element $(\xi, \eta, t) \in \Lambda^{(2)} \times \mathbb{R}$. Hence for any $(\xi_0, \eta_0) \in \Lambda^{(2)}$, there exist sequences $g_i \in \Gamma$ and $t_i \rightarrow +\infty$ such that

$$g_i(\xi, \eta, t + t_i) \rightarrow (\xi_0, \eta_0, 0) \quad \text{as } i \rightarrow \infty.$$

In particular, we have

$$(5.2) \quad g_i(\xi, \eta) \rightarrow (\xi_0, \eta_0) \quad \text{and} \quad \varphi(\beta_\xi^{\theta_1}(g_i^{-1}, e)) \rightarrow -\infty.$$

Since the action of Γ on Λ^{θ_1} is a convergence group action [21, Theorem 4.16], after passing to a subsequence, there exist $a, b \in \Lambda^{\theta_1}$ such that as $i \rightarrow \infty$,

$$g_i|_{\Lambda^{\theta_1} - \{b\}} \rightarrow a \quad \text{uniformly on compact subsets.}$$

That is, for any compact subsets $C_a \subset \Lambda^{\theta_1} - \{a\}$ and $C_b \subset \Lambda^{\theta_1} - \{b\}$,

$$\#\{g_i : g_i C_b \cap C_a \neq \emptyset\} < \infty,$$

or equivalently $\#\{g_i^{-1} : g_i^{-1} C_a \cap C_b \neq \emptyset\} < \infty$. Therefore we have, as $i \rightarrow \infty$,

$$g_i^{-1}|_{\Lambda^{\theta_1} - \{a\}} \rightarrow b \quad \text{uniformly on compact subsets.}$$

Since $g_i(\xi, \eta) \rightarrow (\xi_0, \eta_0)$, we have either

$$(5.3) \quad (a, b) = (\xi_0, \eta) \quad \text{or} \quad (a, b) = (\eta_0, \xi).$$

By the θ_1 -regularity of Γ , we may assume by passing to a subsequence that the sequence $g_i o_1$ (resp. $g_i^{-1} o_1$) converges to some point, say $z \in \Lambda^{\theta_1}$ (resp. $z' \in \Lambda^{\theta_1}$). We claim that $z = a$ and $z' = b$. Write $g_i = k_i b_i \ell_i^{-1} \in KA^+K$ using the Cartan decomposition. By passing to a subsequence, we may assume that $k_i \rightarrow k_0 \in K$ and $\ell_i \rightarrow \ell_0 \in K$. Choose $x \in \Lambda^{\theta_1} - \{\xi, \eta, \xi_0, \eta_0\}$ which is in general position with $\ell_0 w_0 P_{\theta_1}$ and $k_0 P_{\theta_1}$; this is possible by the Zariski density of Γ . Since Γ is θ_1 -regular, we have $\min_{\alpha \in \theta_1} \alpha(\log b_i) \rightarrow \infty$. Hence, by Lemma 2.5, we have

$$g_i x \rightarrow k_0 P_{\theta_1} = z.$$

Since $x \neq b$, we must have $z = a$.

Similarly, the Cartan decomposition $g_i^{-1} = (\ell_i w_0)(w_0^{-1} b_i^{-1} w_0)(w_0^{-1} k_i^{-1}) \in KA^+K$ and the θ_1 -regularity of Γ imply $\min_{\alpha \in \theta_1} \alpha(\log(w_0^{-1} b_i^{-1} w_0)) \rightarrow \infty$. Hence it follows from Lemma 2.5 that

$$g_i^{-1} x \rightarrow \ell_0 w_0 P_{\theta_1} = z'.$$

Since $x \neq a$, we must have $z' = b$, which shows the claim.

Therefore, it suffices to show that $(a, b) = (\eta_0, \xi)$ since we already know that $g_i \xi \rightarrow \xi_0$ and $g_i o_1 \rightarrow a$. Suppose not. Then by (5.3), $(a, b) = (\xi_0, \eta)$, and hence $g_i^{-1} o_1 \rightarrow \eta$. Since $g_i(\xi, \eta) \rightarrow (\xi_0, \eta_0)$ in $\Lambda^{(2)}$, we have $g_i^{-1} o_1 \rightarrow \eta$ conically by Lemma 2.6.

Choose $g \in G_1$ so that $\xi = g P_{\theta_1}$ and $\eta = g w_0 P_{\theta_1}$, noting that we are assuming that $\theta_1 = \mathfrak{i}(\theta_1)$. That g_i^{-1} conically converges to η means that there exist a sequence $k_i \in K$ and a sequence $a_i \rightarrow \infty$ in A^+ such that $\eta = k_i P_{\theta_1}$

for all i and the sequence $g_i k_i a_i$ is bounded. Since $\eta = gw_0 P_{\theta_1} = k_i P_{\theta_1}$, we have for each i , $gw_0 m'_i p_i = k_i$ for some $m'_i \in M_{\theta_1}$ and $p_i \in P$, using $P_{\theta_1} = M_{\theta_1} P$. Since both k_i and m'_i are bounded sequences, the sequence $p_i \in P$ is bounded as well. It implies that the sequence $a_i^{-1} p_i a_i$ is bounded since $a_i \in A^+$. Hence it follows from the boundedness of the sequence $g_i k_i a_i = g_i gw_0 m'_i p_i a_i = g_i gw_0 m'_i a_i (a_i^{-1} p_i a_i)$ that

the sequence $h_i := g_i gw_0 m'_i a_i$ is bounded.

For each i , set $m_i = w_0 m'_i w_0^{-1} \in M_{\theta_1}$. Then

$$\eta = gw_0 P_{\theta_1} = gw_0 m'_i P_{\theta_1} = gm_i w_0 P_{\theta_1}, \quad \xi = gP_{\theta_1} = gm_i P_{\theta_1}$$

and

$$h_i = g_i gw_0 m'_i a_i = g_i gm_i w_0 a_i.$$

Using $\xi = gm_i P_{\theta_1}$, we have

$$\begin{aligned} \beta_{\xi}^{\theta_1}(g_i^{-1}, e) &= \beta_{g_i \xi}^{\theta_1}(e, g_i) = \beta_{g_i \xi}^{\theta_1}(e, h_i) + \beta_{g_i \xi}^{\theta_1}(h_i, g_i) \\ &= \beta_{g_i \xi}^{\theta_1}(e, h_i) + \beta_{\xi}^{\theta_1}(gm_i w_0 a_i, e) \\ &= \beta_{g_i \xi}^{\theta_1}(e, h_i) + \beta_{P_{\theta_1}}^{\theta_1}(w_0 a_i, e) + \beta_{P_{\theta_1}}^{\theta_1}(e, m_i^{-1} g^{-1}). \end{aligned}$$

Since h_i is a bounded sequence, the sequence $\beta_{g_i \xi}^{\theta_1}(e, h_i)$ is bounded by [32, Lemma 5.1]. Similarly, $\beta_{P_{\theta_1}}^{\theta_1}(e, m_i^{-1} g^{-1})$ is bounded. Hence it suffices to show that as $i \rightarrow \infty$,

$$(5.4) \quad \varphi(\beta_{P_{\theta_1}}^{\theta_1}(w_0 a_i, e)) \rightarrow \infty,$$

which yields a contradiction to (5.2). Note that

$$\beta_P(w_0 a_i, e) = \beta_P(w_0 a_i w_0^{-1}, e) = i(\log a_i).$$

Since $h_i = g_i gm_i w_0 a_i$ is bounded and $g_i^{-1} h_i = gm_i w_0 a_i$, we have $\|\mu(g_i^{-1}) - \log a_i\| = \|\mu(g_i) - i(\log a_i)\|$ is uniformly bounded by Lemma 2.1 and the identity (2.1). Therefore

$$\sup_i |\varphi(\mu_{\theta_1}(g_i) - p_{\theta_1}(i(\log a_i)))| < \infty.$$

By the θ_1 -regularity of Γ and the (Γ, θ_1) -properness of φ , $\varphi(\mu_{\theta_1}(g_i)) \rightarrow \infty$ and hence $\varphi(p_{\theta_1}(i(\log a_i))) \rightarrow \infty$ as $i \rightarrow \infty$, which implies (5.4). Hence $\varphi(\beta_{\xi}^{\theta_1}(g_i^{-1}, e)) \rightarrow \infty$, yielding the desired contradiction to (5.2). This shows that $(a, b) = (\eta_0, \xi)$.

Consequently, we have $\xi \in \Lambda_M^{\theta_1}$. Since this holds for $\nu \otimes \nu_i \otimes dt$ -a.e. $(\xi, \eta, t) \in \Lambda^{(2)} \times \mathbb{R}$, we have $\nu(\Lambda_M^{\theta_1}) = 1$.

Myrberg limit set of Γ_{ρ} is ν_{ρ} -full. Note that we have ρ -equivariant continuous maps $f : \Lambda^{\theta_1} \rightarrow \Lambda_{\rho(\Gamma)}^{\theta_2}$ and $f_i : \Lambda^{i(\theta_1)} \rightarrow \Lambda_{\rho(\Gamma)}^{i(\theta_2)}$. As in the previous argument, take $(\xi, \eta, t) \in \Lambda^{(2)} \times \mathbb{R}$ with a dense \mathbb{R}_+ -orbit in Ω_{φ} . Setting $\Lambda_{\rho}^{(2)} := (\Lambda_{\rho}^{\theta} \times \Lambda_{\rho}^{i(\theta)}) \cap \mathcal{F}_{\theta}^{(2)}$, every element of $\Lambda_{\rho}^{(2)}$ is a pair of $(\xi_0, f(\xi_0)) \in \Lambda_{\rho}^{\theta}$

and $(\eta_0, f_1(\eta_0)) \in \Lambda_\rho^{i(\theta)}$ in general position for some $\xi_0, \eta_0 \in \Lambda^{\theta_1}$. Take such elements $\xi_0, \eta_0 \in \Lambda^{\theta_1}$; then $(\xi_0, \eta_0) \in \Lambda^{(2)}$ and hence we again have sequences $g_i \in \Gamma$ and $t_i \rightarrow \infty$ such that $g_i(\xi, \eta, t + t_i) \rightarrow (\xi_0, \eta_0, 0)$ as $i \rightarrow \infty$ as above. As we have shown, it follows that

$$g_i|_{\Lambda^{\theta_1} - \{\xi\}} \rightarrow \eta_0 \quad \text{and} \quad g_i o_1 \rightarrow \eta_0 \quad \text{as } i \rightarrow \infty.$$

For each i , we denote by $\gamma_i = (g_i, \rho(g_i)) \in \Gamma_\rho$. As f and f_1 are ρ -equivariant continuous maps, we have

$$\gamma_i(\xi, f(\xi)) \rightarrow (\xi_0, f(\xi_0)) \quad \text{and} \quad \rho(g_i)|_{\Lambda_{\rho(\Gamma)}^{i(\theta_2)} - \{f_1(\xi)\}} \rightarrow f_1(\eta_0).$$

Hence it remains to show that $\rho(g_i) o_2 \rightarrow f_1(\eta_0)$ as this, together with $g_i o_1 \rightarrow \eta_0$, implies $\gamma_i o \rightarrow (\eta_0, f_1(\eta_0))$.

Since $\rho(\Gamma)$ is θ_2 -regular, it is $i(\theta_2)$ -regular. Write the Cartan decomposition $\rho(g_i) = \tilde{k}_i \tilde{a}_i \tilde{\ell}_i^{-1} \in K A^+ K$. By passing to a subsequence, we may assume that $\tilde{k}_i \rightarrow \tilde{k}_0$ and $\tilde{\ell}_i \rightarrow \tilde{\ell}_0$. Since $\rho(\Gamma)$ is Zariski dense, we can choose $\tilde{x} \in \Lambda_{\rho(\Gamma)}^{i(\theta_2)} - \{f_1(\xi)\}$ which is in general position with $\tilde{\ell}_0 w_0 P_{i(\theta_2)}$. The $i(\theta_2)$ -regularity of $\rho(\Gamma)$ implies that $\min_{\alpha \in i(\theta_2)} \alpha(\log \tilde{a}_i) \rightarrow \infty$. Hence it follows by Lemma 2.5 that

$$\rho(g_i) \tilde{x} \rightarrow \tilde{k}_0 P_{i(\theta_2)} = \lim \rho(g_i) o_2.$$

Since $\tilde{x} \neq f_1(\xi)$, we have $\lim \rho(g_i) o_2 = f_1(\eta_0)$, as desired.

Therefore, we have $(\xi, f(\xi)) \in \Lambda_{\rho, M}^\theta$. Since it holds for $\nu \otimes \nu_i \otimes dt$ -a.e. $(\xi, \eta, t) \in \Lambda^{(2)} \times \mathbb{R}$ and $\nu_\rho = (\text{id} \times f)_* \nu$, we have $\nu_\rho(\Lambda_{\rho, M}^\theta) = 1$, finishing the proof. \square

6. δ -HYPERBOLIC SPACES

In this section, we present certain results on geometric properties of a δ -hyperbolic space; while the qualitative statements in this section is known to experts, it is important for us to express every constant purely in terms of δ . We refer to ([9, Part III], [7], [20], [14, Chapter 1]) for comprehensive expositions.

Let (Z, d_Z) be a proper geodesic metric space. The Gromov product of $y, z \in Z$ with respect to $x \in Z$ is defined as follows:

$$\langle y, z \rangle_x := \frac{1}{2} (d_Z(x, y) + d_Z(x, z) - d_Z(y, z)).$$

It is straightforward to see that for all $x, y, z \in Z$,

$$\langle y, z \rangle_x = \langle z, y \rangle_x \quad \text{and} \quad 0 \leq \langle y, z \rangle_x \leq d_Z(x, y).$$

For $\delta \geq 0$, we call that Z is δ -hyperbolic if

$$\langle w, z \rangle_x \geq \min\{\langle w, y \rangle_x, \langle y, z \rangle_x\} - \delta$$

for all $w, x, y, z \in Z$. The metric space Z is called Gromov hyperbolic if it is δ -hyperbolic for some $\delta \geq 0$.

In the rest of this section, let Z be a proper geodesic δ -hyperbolic space for $\delta \geq 0$. We fix the constant δ and keep track of other constants in terms of δ in the following discussion.

Basic geometry. We first discuss some basic geometry of Z . The following standard lemma says that the Gromov product gives the length of the initial segments of two geodesics from a common point with uniformly bounded Hausdorff distance:

Lemma 6.1. *Let $x, y, z \in Z$ and fix geodesic segments $[x, y], [x, z] \subset Z$ between x and y , and x and z , respectively. If $y' \in [x, y]$ and $z' \in [x, z]$ are such that $d_Z(x, y') = d_Z(x, z') \leq \langle y, z \rangle_x$, then $d_Z(y', z') \leq 4\delta$.*

Proof. Since Z is δ -hyperbolic, we have

$$\begin{aligned} \langle y', z' \rangle_x &\geq \min\{\langle y', z \rangle_x, \langle z, z' \rangle_x\} - \delta \\ &\geq \min\{\min\{\langle y', y \rangle_x, \langle y, z \rangle_x\} - \delta, \langle z, z' \rangle_x\} - \delta \\ &= \min\{\min\{d_Z(x, y'), \langle y, z \rangle_x\} - \delta, d_Z(x, z')\} - \delta \\ &= d_Z(x, y') - 2\delta. \end{aligned}$$

On the other hand, $\langle y', z' \rangle_x = d_Z(x, y') - \frac{1}{2}d_Z(y', z')$, from which the claim follows. \square

As a corollary, we deduce that every geodesic triangle in Z is uniformly thin:

Corollary 6.2. *Let $x, y, z \in Z$ and fix geodesic segments $[x, y], [y, z], [x, z] \subset Z$ between x and y , y and z , and x and z , respectively. Then $[x, y]$ is contained in the 4δ -neighborhood of $[x, z] \cup [y, z]$.*

We also obtain the interpretation that the Gromov product roughly measures a distance between a point and a geodesic segment.

Corollary 6.3. *Let $x, y, z \in Z$ and fix a geodesic segment $[y, z] \subset Z$ between y and z . Then*

$$d_Z(x, [y, z]) - 4\delta \leq \langle y, z \rangle_x \leq d_Z(x, [y, z]).$$

Proof. Let $w \in [y, z]$ be such that $d_Z(x, w) = d_Z(x, [y, z])$. Then

$$\begin{aligned} \langle y, z \rangle_x &= \frac{1}{2}(d_Z(x, y) + d_Z(x, z) - d_Z(y, z)) \\ &= \frac{1}{2}(d_Z(x, y) - d_Z(y, w)) + \frac{1}{2}(d_Z(x, z) - d_Z(w, z)) \\ &\leq d_Z(x, w) = d_Z(x, [y, z]). \end{aligned}$$

To see the lower bound, fix a geodesic segment $[x, y] \subset Z$ between x and y and let $y' \in [x, y]$ be the point such that $d_Z(x, y') = \langle y, z \rangle_x$. Since $d_Z(x, y) = \langle y, z \rangle_x + \langle x, z \rangle_y$, we have $d_Z(y, y') = \langle x, z \rangle_y$. Let $z' \in [y, z]$ be the point such that $d_Z(y, z') = \langle x, z \rangle_y$. By Lemma 6.1, $d_Z(y', z') \leq 4\delta$, and hence

$$d_Z(x, z') \leq d_Z(x, y') + d_Z(y', z') \leq \langle y, z \rangle_x + 4\delta.$$

Since $d_Z(x, [y, z]) \leq d_Z(x, z')$, this finishes the proof. \square

Note that in Corollary 6.3, the choice of a geodesic segment was made while the Gromov product does not involve any choice of a geodesic segment. Indeed, geodesics between two points are stable:

Corollary 6.4. *Let $y, z \in Z$. Then two geodesic segments in Z between y and z have Hausdorff distance at most 4δ .*

Proof. Let $\sigma_1, \sigma_2 \subset Z$ be two geodesics between y and z . Fix any $x \in \sigma_1$. Then $\langle y, z \rangle_x = 0$. By Corollary 6.3, this implies $d_Z(x, \sigma_1) \leq 4\delta$. Since x is arbitrary, σ_2 is contained in the 4δ -neighborhood of σ_1 . The same argument switching σ_1 and σ_2 finishes the proof. \square

The following will be a useful observation:

Corollary 6.5. *Let $x \in Z$ and σ be a geodesic segment in Z . Let $y \in \sigma$ be such that $d_Z(x, y) = d_Z(x, \sigma)$. Then for any $z \in \sigma$, we have*

$$d_Z(x, y) + d_Z(y, z) - 8\delta \leq d_Z(x, z) \leq d_Z(x, y) + d_Z(y, z).$$

Proof. The upper bound is straightforward. By Corollary 6.3, we have

$$\begin{aligned} d_Z(x, y) &\leq \langle y, z \rangle_x + 4\delta \\ &= \frac{1}{2}(d_Z(x, y) + d_Z(x, z) - d_Z(y, z)) + 4\delta. \end{aligned}$$

This implies the lower bound. \square

Gromov boundary. An isometric embedding $\sigma : [0, \infty) \rightarrow Z$, or its image in Z , is called a geodesic ray in Z . The Gromov boundary of Z is defined as the set of all equivalence classes of geodesic rays in Z :

$$\partial Z := \{\sigma : [0, \infty) \rightarrow Z, \text{ a geodesic ray}\} / \sim$$

where $\sigma \sim \sigma'$ if the Hausdorff distance between two geodesic rays $\sigma([0, \infty))$ and $\sigma'([0, \infty))$ is finite. We denote by $\sigma(\infty) \in \partial Z$ the equivalence class of the geodesic ray $\sigma : [0, \infty) \rightarrow Z$. Fixing a basepoint in Z , the Gromov boundary ∂Z is visible from the basepoint:

Lemma 6.6. [9, Lemma III.3.1] *Let $x \in Z$ and $\xi \in \partial Z$. Then there exists a geodesic ray $\sigma : [0, \infty) \rightarrow Z$ such that $\sigma(0) = x$ and $\sigma(\infty) = \xi$.*

Moreover, this visualization is stable under the choice of the basepoint:

Lemma 6.7. [9, Lemma III.3.3] *Let $\sigma_1, \sigma_2 : [0, \infty) \rightarrow Z$ be geodesic rays with $\sigma_1(\infty) = \sigma_2(\infty)$.*

- (1) *If $\sigma_1(0) = \sigma_2(0)$, then $d_Z(\sigma_1(t), \sigma_2(t)) \leq 8\delta$ for all $t \geq 0$.*
- (2) *In general, there exist $T_1, T_2 \geq 0$ such that $d_Z(\sigma_1(t+T_1), \sigma_2(t+T_2)) \leq 20\delta$ for all $t \geq 0$.*

Hence, fixing a basepoint $x \in Z$, we can identify the Gromov boundary of Z with the set of all equivalence classes of geodesic rays in Z based at x :

$$\partial Z = \{\sigma : [0, \infty) \rightarrow Z, \text{ a geodesic ray with } \sigma(0) = x\} / \sim$$

where $\sigma \sim \sigma'$ if $\sigma(t) \sim \sigma'(t) \leq 8\delta$ for all $t \geq 0$. Under this identification, a natural topology on ∂Z is given as follows: for $\xi \in \partial Z$ and $r \geq 0$, we set

$$V(\xi, r) := \left\{ \eta \in \partial Z : \begin{array}{l} \text{for some geodesic rays } \sigma, \sigma' \text{ from } x \\ \text{with } \sigma(\infty) = \xi \text{ and } \sigma'(\infty) = \eta, \\ \text{we have } \liminf_{t \rightarrow \infty} \langle \sigma(t), \sigma'(t) \rangle_x \geq r \end{array} \right\}$$

which consists of the geodesic rays from x that are 8δ -close to σ for a long time. We topologize ∂Z by setting $\{V(\xi, r) : \xi \in \partial Z, r \geq 0\}$ to be the basis.

We now consider $\bar{Z} = Z \cup \partial Z$ and give it a natural topology. We say that a sequence (x_i) converges to infinity if $\liminf_{i, j \rightarrow \infty} \langle x_i, x_j \rangle_x = \infty$. To topologize \bar{Z} , it is useful to associate a geodesic ray $\sigma : [0, \infty) \rightarrow Z$ with a sequence $(\sigma(i))_{i \in \mathbb{N}}$ which converges to infinity. This gives a map

$$\partial Z \rightarrow \{(x_i) \subset Z, \text{ a sequence converging to infinity}\} / \sim$$

where $(x_i) \sim (y_i)$ if $\liminf_{i, j \rightarrow \infty} \langle x_i, y_j \rangle_x = \infty$. The above map is indeed a bijection, and hence we identify them as well. We denote by $[(x_i)]$ the equivalence class of the sequence (x_i) . Similar to the above, for $\xi \in \partial Z$ and $r \geq 0$, we set

$$U(\xi, r) := \left\{ \eta \in \partial Z : \begin{array}{l} \text{for some sequences } (x_i), (y_i) \\ \text{with } [(x_i)] = \xi \text{ and } [(y_i)] = \eta, \\ \text{we have } \liminf_{i, j \rightarrow \infty} \langle x_i, y_j \rangle_x \geq r \end{array} \right\}.$$

The topology on ∂Z given by setting $\{U(\xi, r) : \xi \in \partial Z, r \geq 0\}$ as a basis is equivalent to the one defined in terms of $V(\xi, r)$. To obtain a basis for \bar{Z} , we also consider for $\xi \in \partial Z$ and $r \geq 0$

$$U'(\xi, r) := U(\xi, r) \cup \left\{ y \in Z : \begin{array}{l} \text{for some sequence } (x_i) \text{ with } [(x_i)] = \xi, \\ \text{we have } \liminf_{i \rightarrow \infty} \langle x_i, y \rangle_x \geq r \end{array} \right\}.$$

Then setting $\{U'(\xi, r) : \xi \in \partial Z, r \geq 0\}$ and metric balls in Z to be the basis, \bar{Z} is equipped with the topology. In this topology, a sequence x_i in Z converges to $\xi \in \partial Z$ if and only if $\xi = [(x_i)]$. The spaces ∂Z and \bar{Z} equipped with these topologies are compact, and the topologies do not depend on the choice of the basepoint x . We refer to ([9], [20]) for details.

Extended Gromov product. We extend the notion of the Gromov product to \bar{Z} : for $y, z \in \bar{Z}$ and $x \in Z$, the Gromov product of y and z with respect to x is defined as

$$\langle y, z \rangle_x := \sup \liminf_{i, j \rightarrow \infty} \langle y_i, z_j \rangle_x$$

where the supremum is taken over all sequences (y_i) and (z_j) in Z such that $y = \lim_i y_i$ and $z = \lim_j z_j$. We note the following properties of the extended Gromov product:

Lemma 6.8. [9, Remark III.3.17] *Fix $x \in Z$.*

- (1) *For $y, z \in \partial Z$, $\langle y, z \rangle_x = \infty$ if and only if $y = z$.*
- (2) *For $w, y, z \in \bar{Z}$, we have*

$$\langle w, z \rangle_x \geq \min\{\langle w, y \rangle_x, \langle y, z \rangle_x\} - 2\delta.$$

- (3) *For $y, z \in \bar{Z}$ and sequences $(y_i), (z_j)$ in Z with $\lim_i y_i = y$ and $\lim_j z_j = z$, we have*

$$\langle y, z \rangle_x - 2\delta \leq \liminf_{i,j} \langle y_i, z_j \rangle_x \leq \langle y, z \rangle_x.$$

- (4) *For $z \in \partial Z$ and a sequence (z_i) in ∂Z , $z_i \rightarrow z$ as $i \rightarrow \infty$ if and only if $\langle z, z_i \rangle_x \rightarrow \infty$ as $i \rightarrow \infty$.*

Given two distinct points $y, z \in \partial Z$, there exists a bi-infinite geodesic $\sigma : \mathbb{R} \rightarrow Z$ connecting y and z , i.e., $\sigma(-\infty) = y$ and $\sigma(\infty) = z$ [9, Lemma III.3.2]. Hence in general we can consider a geodesic between two points in \bar{Z} . The extended Gromov product $\langle y, z \rangle_x$ also measures the crude distance from x to a geodesic between $y, z \in \bar{Z}$.

Corollary 6.9. *Let $x \in Z$ and $y, z \in \bar{Z}$ be distinct points. Let $[y, z]$ be a geodesic connecting y and z . Then we have*

$$d_Z(x, [y, z]) - 4\delta \leq \langle y, z \rangle_x \leq d_Z(x, [y, z]) + 2\delta.$$

Proof. Let $w \in [y, z]$ be such that $d_Z(x, w) = d_Z(x, [y, z])$. Let (y_i) and (z_j) be sequences of points on $[y, z]$ such that $\lim_i y_i = y$ and $\lim_j z_j = z$. For each i and j , let $[y_i, z_j] \subset [y, z]$ be the segment between y_i and z_j . Then for large enough i and j , we have $w \in [y_i, z_j]$ and hence by Corollary 6.3,

$$d_Z(x, [y, z]) - 4\delta \leq \langle y_i, z_j \rangle_x \leq d_Z(x, [y, z])$$

since $d_Z(x, [y_i, z_j]) = d_Z(x, [y, z])$. Applying Lemma 6.8(3) finishes the proof. \square

As in Corollary 6.4, we also obtain the stability of geodesics between two points in \bar{Z} .

Corollary 6.10. *Let $y, z \in \bar{Z}$. Then two geodesics between y and z have Hausdorff distance at most 6δ .*

Proof. Suppose first that $y, z \in \partial Z$. Let $\sigma_1, \sigma_2 : \mathbb{R} \rightarrow Z$ be two bi-infinite geodesics between y and z . Let $x \in \sigma_1(\mathbb{R})$. Then by Corollary 6.9, we have

$$d_Z(x, \sigma_2(\mathbb{R})) \leq \langle y, z \rangle_x + 4\delta.$$

On the other hand, $0 = \liminf_{t \rightarrow \infty} \langle \sigma_1(-t), \sigma_1(t) \rangle_x \geq \langle y, z \rangle_x - 2\delta$ by Lemma 6.8(3). Therefore we have

$$d_Z(x, \sigma_2(\mathbb{R})) \leq 6\delta.$$

Since x is arbitrary, this finishes the proof in this case. The case when one of y and z is in Z can be handled similarly. \square

Visual metric. Indeed, ∂Z can be equipped with a natural metric-like function. From the above observation, it is natural to consider the following function which plays a role of metric on ∂Z , which we call the visual metric on ∂Z , although it may not satisfy the triangle inequality in general:

Definition 6.11. Let $x \in Z$. We define a function $d_x : \partial Z \times \partial Z \rightarrow \mathbb{R}$ as

$$d_x(y, z) := e^{-2\langle y, z \rangle_x}.$$

For $y \in \partial Z$ and $r > 0$ we consider the d_x -ball

$$B_x(y, r) := \{z \in \partial Z : d_x(y, z) < r\}.$$

Usually the visual metric is defined without the multiplication by 2. However, we defined it as above in order to simplify the later computation. The visual metric is compatible to a genuine metric on ∂Z after taking a suitable power:

Proposition 6.12. [9, Proposition III.3.21] *Let $x \in Z$. For any small enough $\varepsilon > 0$, there exists a constant c_ε and a metric d_ε on ∂Z such that*

$$d_\varepsilon(y, z) \leq d_x(y, z)^\varepsilon \leq c_\varepsilon d_\varepsilon(y, z)$$

for all $y, z \in \partial Z$.

It follows from Lemma 6.8(2) that for any $w, y, z \in \partial Z$, we have

$$(6.1) \quad d_x(w, z) \leq e^{4\delta}(d_x(w, y) + d_x(y, z))$$

From this we deduce the following Vitali-type covering lemma:

Lemma 6.13. *Let $x \in Z$ and $B_x(y_1, r_1), \dots, B_x(y_n, r_n)$ a finite collection of d_x -balls for $y_i \in \partial Z$ and $r_i > 0$. Then there a subcollection of disjoint balls $B_x(y_{i_1}, r_{i_1}), \dots, B_x(y_{i_k}, r_{i_k})$ such that*

$$\bigcup_{i=1}^n B_x(y_i, r_i) \subset \bigcup_{j=1}^k B_x(y_{i_j}, 3e^{8\delta}r_{i_j}).$$

Proof. Given a finite collection $B_x(y_1, r_1), \dots, B_x(y_n, r_n)$ of d_x -balls, we rearrange them so that we may assume $r_1 \geq \dots \geq r_n$. Let $i_1 = 1$ and for each $j \geq 2$, we set $i_j = \min\{i > i_{j-1} : B_x(y_i, r_i) \cap \bigcup_{\ell=1}^{i_j-1} B_x(y_\ell, r_\ell) = \emptyset\}$. Then we obtain a subcollection $B_x(y_{i_1}, r_{i_1}), \dots, B_x(y_{i_k}, r_{i_k})$ consisting of disjoint balls.

For each i , $B_x(y_i, r_i)$ intersects $B_x(y_{i_j}, r_{i_j})$ for some j such that $r_{i_j} \geq r_i$. Choosing a point $y \in B_x(y_i, r_i) \cap B_x(y_{i_j}, r_{i_j})$, it follows from (6.1) that for any $z \in B_x(y_i, r_i)$,

$$\begin{aligned} d_x(z, y_{i_j}) &\leq e^{4\delta}(d_x(z, y_i) + d_x(y_i, y_{i_j})) \\ &\leq e^{4\delta}(r_i + e^{4\delta}(d_x(y_i, y) + d_x(y, y_{i_j}))) \\ &\leq e^{4\delta}(r_i + e^{4\delta}(r_i + r_{i_j})) \leq 3e^{8\delta}r_{i_j}. \end{aligned}$$

Hence $B_x(y_i, r_i) \subset B_x(y_{i_j}, 3e^{8\delta}r_{i_j})$. This finishes the proof. \square

Busemann functions. Let $\sigma : [0, \infty) \rightarrow Z$ be a geodesic ray and $y, z \in Z$. Then the following limit is well-defined and satisfies the following inequality:

$$(6.2) \quad -d_Z(y, z) \leq \lim_{t \rightarrow \infty} d_Z(y, \sigma(t)) - d_Z(z, \sigma(t)) \leq d_Z(y, z).$$

Therefore, we define the Busemann function as follows:

$$\beta_\sigma(y, z) := \lim_{t \rightarrow \infty} d_Z(y, \sigma(t)) - d_Z(z, \sigma(t)).$$

Observe that: for $w, y, z \in Z$,

- (1) we have $|\beta_\sigma(y, z)| \leq d_Z(y, z)$;
- (2) we have $\beta_\sigma(y, z) = -\beta_\sigma(z, y)$;
- (3) we have $\beta_\sigma(w, z) = \beta_\sigma(w, y) + \beta_\sigma(y, z)$.

The Busemann function depends only on the endpoint at ∂Z , independent of a choice of a geodesic ray, up to a uniform error.

Lemma 6.14. *Let $\sigma, \sigma' : [0, \infty) \rightarrow Z$ be geodesic rays such that $\sigma(\infty) = \sigma'(\infty)$. Then for any $y, z \in Z$, we have*

$$|\beta_\sigma(y, z) - \beta_{\sigma'}(y, z)| \leq 40\delta.$$

Proof. By Lemma 6.7, there exists $T, T' > 0$ such that

$$d_Z(\sigma(t+T), \sigma'(t+T')) \leq 20\delta$$

for all $t \geq 0$. This implies the desired inequality. \square

Moreover, the Busemann function is stable under the change of the endpoint.

Lemma 6.15. *Let $x \in Z$ and $r > 0$. Let $y, z \in Z$ be such that $d_Z(x, y) < r - 10\delta$ and $d_Z(x, z) < r - 10\delta$. Let $\sigma, \sigma' : [0, \infty) \rightarrow Z$ be geodesic rays with $\langle \sigma(\infty), \sigma'(\infty) \rangle_x > r$. If $\sigma(0) = \sigma'(0) = x$, then*

$$|\beta_\sigma(y, z) - \beta_{\sigma'}(y, z)| \leq 72\delta.$$

In general,

$$|\beta_\sigma(y, z) - \beta_{\sigma'}(y, z)| \leq 152\delta.$$

Proof. Suppose first that $\sigma(0) = \sigma'(0) = x$. By Lemma 6.8(3), we have for all large $t > 0$ that

$$\langle \sigma(t), \sigma'(t) \rangle_x > r - 2\delta.$$

Fix such $t > 0$. Let $t_0 \geq 0$ be such that $d_Z(y, \sigma([0, \infty))) = d_Z(y, \sigma(t_0))$. By Corollary 6.5, we have

$$(6.3) \quad t_0 + d_Z(\sigma(t_0), y) - 8\delta \leq d_Z(x, y) \leq t_0 + d_Z(\sigma(t_0), y).$$

In particular, we have $t_0 \leq d_Z(x, y) + 8\delta < r - 2\delta < \langle \sigma(t), \sigma'(t) \rangle_x$. Hence by Lemma 6.1, we have

$$(6.4) \quad d_Z(\sigma(t_0), \sigma'(t_0)) \leq 4\delta.$$

Similarly, letting $t'_0 \geq 0$ be such that $d_Z(y, \sigma'([0, \infty))) = d_Z(y, \sigma'(t'_0))$, it follows from Corollary 6.5 that

$$(6.5) \quad t'_0 + d_Z(\sigma'(t'_0), y) - 8\delta \leq d_Z(x, y) \leq t'_0 + d_Z(\sigma'(t'_0), y).$$

Similarly, by Lemma 6.1, we have

$$(6.6) \quad d_Z(\sigma(t'_0), \sigma'(t'_0)) \leq 4\delta.$$

Combining (6.3) and (6.5), we have

$$d_Z(\sigma'(t'_0), y) - d_Z(\sigma(t_0), y) - 8\delta \leq t_0 - t'_0 \leq d_Z(\sigma'(t'_0), y) - d_Z(\sigma(t_0), y) + 8\delta.$$

Since $d_Z(y, \sigma'([0, \infty)) = d_Z(y, \sigma'(t'_0))$, we have

$$\begin{aligned} t_0 - t'_0 &\leq d_Z(\sigma'(t'_0), y) - d_Z(\sigma(t_0), y) + 8\delta \\ &\leq d_Z(\sigma'(t_0), y) - d_Z(\sigma(t_0), y) + 8\delta \\ &\leq d_Z(\sigma'(t_0), \sigma(t_0)) + 8\delta \\ &\leq 12\delta \end{aligned}$$

where the last inequality is by (6.4). Similarly, we have

$$\begin{aligned} t_0 - t'_0 &\geq d_Z(\sigma'(t'_0), y) - d_Z(\sigma(t_0), y) - 8\delta \\ &\geq d_Z(\sigma'(t'_0), y) - d_Z(\sigma(t'_0), y) - 8\delta \\ &\leq -d_Z(\sigma'(t'_0), \sigma(t'_0)) - 8\delta \\ &\leq -12\delta \end{aligned}$$

where the last inequality is by (6.6). Therefore we obtain

$$|t_0 - t'_0| \leq 12\delta,$$

and hence

$$d_Z(\sigma(t_0), \sigma'(t'_0)) \leq d_Z(\sigma(t_0), \sigma(t'_0)) + d_Z(\sigma(t'_0), \sigma'(t'_0)) \leq 16\delta.$$

Now for all large $t > 0$, it follows from Corollary 6.5 that

$$\begin{aligned} d_Z(x, \sigma(t)) - d_Z(y, \sigma(t_0)) - d_Z(\sigma(t_0), \sigma(t)) \\ \leq d_Z(x, \sigma(t)) - d_Z(y, \sigma(t)) \\ \leq d_Z(x, \sigma(t)) - d_Z(y, \sigma(t_0)) - d_Z(\sigma(t_0), \sigma(t)) + 8\delta. \end{aligned}$$

This implies

$$t_0 - d_Z(y, \sigma(t_0)) \leq d_Z(x, \sigma(t)) - d_Z(y, \sigma(t)) \leq t_0 - d_Z(y, \sigma(t_0)) + 8\delta$$

for all large enough $t > 0$, and therefore we have

$$t_0 - d_Z(y, \sigma(t_0)) \leq \beta_\sigma(x, y) \leq t_0 - d_Z(y, \sigma(t_0)) + 8\delta.$$

Similarly, we also have

$$t'_0 - d_Z(y, \sigma'(t'_0)) \leq \beta_{\sigma'}(x, y) \leq t'_0 - d_Z(y, \sigma'(t'_0)) + 8\delta.$$

Hence, we have

$$\begin{aligned} |\beta_\sigma(x, y) - \beta_{\sigma'}(x, y)| &\leq |t_0 - t'_0| + |d_Z(y, \sigma(t_0)) - d_Z(y, \sigma'(t'_0))| + 8\delta \\ &\leq 12\delta + 16\delta + 8\delta = 36\delta. \end{aligned}$$

By the same argument replacing y with z , we also have

$$|\beta_\sigma(x, z) - \beta_{\sigma'}(x, z)| \leq 36\delta.$$

Therefore, it follows that

$$|\beta_\sigma(y, z) - \beta_{\sigma'}(y, z)| \leq 72\delta,$$

proving the first claim.

The last claim follows from the first claim, by applying Lemma 6.6 and Lemma 6.14. \square

Lemma 6.16. *Let $\sigma_1, \sigma_2 : [0, \infty) \rightarrow Z$ be geodesic rays from $x \in Z$. For any $w \in Z$ on a bi-infinite geodesic between $\sigma_1(\infty)$ and $\sigma_2(\infty)$, we have*

$$\langle \sigma_1(\infty), \sigma_2(\infty) \rangle_x - 42\delta \leq \frac{1}{2} (\beta_{\sigma_1}(x, w) + \beta_{\sigma_2}(x, w)) \leq \langle \sigma_1(\infty), \sigma_2(\infty) \rangle_x.$$

Proof. Let $[\sigma_1(\infty), \sigma_2(\infty)]$ be a bi-infinite geodesic in Z between $\sigma_1(\infty)$ and $\sigma_2(\infty)$ and $w \in [\sigma_1(\infty), \sigma_2(\infty)]$. We then have

$$\begin{aligned} & \beta_{\sigma_1}(x, w) + \beta_{\sigma_2}(x, w) \\ &= \lim_{t \rightarrow \infty} d_Z(x, \sigma_1(t)) - d_Z(w, \sigma_1(t)) + d_Z(x, \sigma_2(t)) - d_Z(w, \sigma_2(t)) \\ &\leq \liminf_{t \rightarrow \infty} d_Z(x, \sigma_1(t)) + d_Z(x, \sigma_2(t)) - d_Z(\sigma_1(t), \sigma_2(t)) \\ &= 2 \liminf_{t \rightarrow \infty} \langle \sigma_1(t), \sigma_2(t) \rangle_x. \end{aligned}$$

Hence, by Lemma 6.8(3), the upper bound follows.

To see the lower bound, let $\sigma'_1, \sigma'_2 : [0, \infty) \rightarrow Z$ be geodesic rays that parametrize the segments of $[\sigma_1(\infty), \sigma_2(\infty)]$ from w to $\sigma_1(\infty)$, from w to $\sigma_2(\infty)$, respectively. Then

$$\begin{aligned} & \beta_{\sigma'_1}(x, w) + \beta_{\sigma'_2}(x, w) \\ &= \lim_{t \rightarrow \infty} d_Z(x, \sigma'_1(t)) - d_Z(w, \sigma'_1(t)) + d_Z(x, \sigma'_2(t)) - d_Z(w, \sigma'_2(t)) \\ &= \lim_{t \rightarrow \infty} d_Z(x, \sigma'_1(t)) + d_Z(x, \sigma'_2(t)) - d_Z(\sigma'_1(t), \sigma'_2(t)) \\ &= 2 \lim_{t \rightarrow \infty} \langle \sigma'_1(t), \sigma'_2(t) \rangle_x. \end{aligned}$$

Therefore, by Lemma 6.14 and Lemma 6.8(3), we have

$$\begin{aligned} \beta_{\sigma_1}(x, w) + \beta_{\sigma_2}(x, w) &\geq \beta_{\sigma'_1}(x, w) + \beta_{\sigma'_2}(x, w) - 80\delta \\ &= 2 \lim_{t \rightarrow \infty} \langle \sigma'_1(t), \sigma'_2(t) \rangle_x - 80\delta \\ &\geq 2 \langle \sigma_1(\infty), \sigma_2(\infty) \rangle_x - 84\delta. \end{aligned}$$

This implies the desired lower bound. \square

We can now compare two visual metrics in terms of Busemann functions:

Lemma 6.17. *Let $y, z \in \partial Z$ and $x, x' \in Z$. Let $\sigma'_1, \sigma'_2 : [0, \infty) \rightarrow Z$ be geodesic rays from x' to y' and z' respectively. Let $\sigma_1, \sigma_2 : [0, \infty) \rightarrow Z$ be any geodesic rays with $\sigma_1(\infty) = y$ and $\sigma_2(\infty) = z$. Then*

$$\begin{aligned} d_{x'}(y, z) &\leq e^{164\delta} e^{\beta_{\sigma'_1}(x, x') + \beta_{\sigma'_2}(x, x')} d_x(y, z) \\ &\leq e^{244\delta} e^{\beta_{\sigma_1}(x, x') + \beta_{\sigma_2}(x, x')} d_x(y, z). \end{aligned}$$

Proof. Let $[y, z]$ be a bi-infinite geodesic between y and z and let $w \in [y, z]$. By Lemma 6.16 and Lemma 6.14, we have

$$\begin{aligned} 2\langle y, z \rangle_{x'} &\geq \beta_{\sigma'_1}(x', w) + \beta_{\sigma'_2}(x', w) \\ &= \beta_{\sigma'_1}(x, w) + \beta_{\sigma'_2}(x, w) + \beta_{\sigma'_1}(x', x) + \beta_{\sigma'_2}(x', x) \\ &\geq \beta_{\sigma_1}(x, w) + \beta_{\sigma_2}(x, w) + \beta_{\sigma'_1}(x', x) + \beta_{\sigma'_2}(x', x) - 80\delta \\ &\geq 2\langle y, z \rangle_x + \beta_{\sigma'_1}(x', x) + \beta_{\sigma'_2}(x', x) - 164\delta \end{aligned}$$

Therefore,

$$d_{x'}(y, z) \leq e^{164\delta} e^{\beta_{\sigma'_1}(x, x') + \beta_{\sigma'_2}(x, x')} d_x(y, z).$$

Applying Lemma 6.14 again, we deduce

$$e^{164\delta} e^{\beta_{\sigma'_1}(x, x') + \beta_{\sigma'_2}(x, x')} d_x(y, z) \leq e^{244\delta} e^{\beta_{\sigma_1}(x, x') + \beta_{\sigma_2}(x, x')} d_x(y, z).$$

□

Shadows. Let $x \in Z$, $y \in \bar{Z}$, and $R > 0$. The shadow of R -ball at x viewed from y is defined as

$$O_R^Z(y, x) = \{z \in \partial Z : \langle y, z \rangle_x < R\}.$$

Busemann function is comparable to the distance in a shadow:

Lemma 6.18. *Let $x, y \in Z$ and $R > 0$. Let $\sigma : [0, \infty) \rightarrow Z$ be a geodesic ray such that $\sigma(\infty) \in O_R^Z(y, x)$. Then*

$$|\beta_\sigma(y, x) - d_Z(y, x)| < 2R + 48\delta.$$

Proof. Fix any geodesic $\sigma' : [0, \infty) \rightarrow Z$ from y to $\sigma(\infty)$. By Corollary 6.9, there exists $t_0 \geq 0$ such that $d_Z(x, \sigma'(t_0)) < R + 4\delta$. We then have

$$\begin{aligned} |\beta_{\sigma'}(y, x) - d_Z(y, \sigma'(t_0))| &= \left| \lim_{t \rightarrow \infty} d_Z(\sigma'(t_0), \sigma'(t)) - d_Z(x, \sigma'(t)) \right| \\ &\leq d_Z(x, \sigma'(t_0)) < R + 4\delta. \end{aligned}$$

By Lemma 6.14, we have

$$\begin{aligned} |\beta_\sigma(y, x) - d_Z(y, x)| &\leq 40\delta + |\beta_{\sigma'}(y, x) - d_Z(y, \sigma'(t_0))| \\ &\quad + |d_Z(y, \sigma'(t_0)) - d_Z(y, x)| \\ &< 40\delta + (R + 4\delta) + (R + 4\delta) = 2R + 48\delta. \end{aligned}$$

□

Shadows viewed from ∂Z can be approximated by shadows viewed from Z .

Lemma 6.19. *Let $x \in Z$ and (y_i) be a sequence in Z such that $\lim_i y_i = y \in \partial Z$. Then for any $R > 0$, we have*

$$O_R^Z(y, x) \subset O_{R+2\delta}^Z(y_i, x)$$

for all $i \geq 1$ large enough.

Proof. Let $z \in O_R^Z(y, x)$. Then by Lemma 6.8(2), we have for each $i \geq 1$ that

$$R > \langle y, z \rangle_x \geq \min\{\langle y, y_i \rangle_x, \langle y_i, z \rangle_x\} - 2\delta.$$

Since $y_i \rightarrow y$ as $i \rightarrow \infty$, for i large enough so that $\langle y, y_i \rangle_x > R + 2\delta$, we have

$$\langle y_i, z \rangle_x < R + 2\delta.$$

This shows the claim. \square

Isometries. Let $g \in \text{Isom}(Z)$ be an isometry of Z . Then $g : Z \rightarrow Z$ extends to a homeomorphism $g : \bar{Z} \rightarrow \bar{Z}$. One can see that for $x, w \in Z$, $y, z \in \bar{Z}$, and a geodesic ray $\sigma : [0, \infty) \rightarrow Z$,

$$\langle gy, gz \rangle_{gx} = \langle y, z \rangle_x \quad \text{and} \quad \beta_{g\sigma}(gx, gw) = \beta_\sigma(x, w).$$

Isometries of Z are classified into three categories. Let $g \in \text{Isom}(Z)$. Then either one of the following holds:

- (1) g is elliptic, i.e., $\{g^n x : n \in \mathbb{Z}\}$ is bounded for any $x \in Z$;
- (2) g is parabolic, i.e., g is not elliptic and has exactly one fixed point in ∂Z ; or
- (3) g is loxodromic, i.e., g is not elliptic and has exactly two fixed points in ∂Z .

If $g \in \text{Isom}(X)$ is loxodromic, we can denote two fixed points by $y_g, y_{g^{-1}} \in \partial Z$ so that $g^n x \rightarrow y_g$ as $n \rightarrow \infty$ for all $x \neq y_{g^{-1}}$ and $g^{-n} x \rightarrow y_{g^{-1}}$ as $n \rightarrow \infty$ for all $x \neq y_g$. We call y_g and $y_{g^{-1}}$ the attracting and repelling fixed points of g respectively.

For $g \in \text{Isom}(X)$, we define its asymptotic translation length by

$$\ell(g) := \lim_{n \rightarrow \infty} \frac{d_Z(x, g^n x)}{n}$$

for $x \in Z$. This does not depend on the choice of x . It is clear that $\ell(hgh^{-1}) = \ell(g)$ and $\ell(g^n) = |n|\ell(g)$ for all $g, h \in \text{Isom}(X)$ and $n \in \mathbb{Z}$.

Lemma 6.20. *Let $g \in \text{Isom}(X)$ be loxodromic and $[y_{g^{-1}}, y_g]$ a geodesic between $y_{g^{-1}}$ and y_g . Let $x \in [y_{g^{-1}}, y_g]$ and $\sigma_0 : [0, \infty) \rightarrow [y_{g^{-1}}, y_g]$ be a parametrization from x to y_g . Then*

$$|\beta_{\sigma_0}(x, gx) - \ell(g)| \leq 12\delta.$$

Moreover, if $\sigma : [0, \infty) \rightarrow Z$ is a geodesic ray with $\sigma(\infty) = y_g$ and $w \in Z$, then

$$|\beta_\sigma(w, gw) - \ell(g)| \leq 92\delta.$$

Proof. Note that $g^n x \rightarrow y_g$ as $n \rightarrow \infty$ and $g^n x$ always belongs to a geodesic $g^n[y_{g^{-1}}, y_g]$ between $y_{g^{-1}}$ and y_g . Hence for all large n , there exists $t_n \geq 0$ such that $d_Z(g^n x, \sigma_0(t_n)) \leq 6\delta$ by Lemma 6.10. We then have for each $n \geq 1$

that

$$\begin{aligned}
 & |\beta_{\sigma_0}(x, g^n x) - d_Z(x, \sigma_0(t_n))| \\
 &= \lim_{t \rightarrow \infty} |d_Z(x, \sigma_0(t)) - d_Z(x, \sigma_0(t_n)) - d_Z(g^n x, \sigma_0(t))| \\
 &= \lim_{t \rightarrow \infty} |d_Z(\sigma_0(t_n), \sigma_0(t)) - d_Z(g^n x, \sigma_0(t))| \leq 6\delta.
 \end{aligned}$$

Together with $d_Z(g^n x, \sigma_0(t_n)) \leq 6\delta$, this implies

$$|\beta_{\sigma_0}(x, g^n x) - d_Z(x, g^n x)| \leq 12\delta.$$

On the other hand, we have

$$\beta_{\sigma_0}(x, g^n x) = \sum_{i=0}^{n-1} \beta_{\sigma_0}(g^i x, g^{i+1} x) = \sum_{i=0}^{n-1} \beta_{g^{-i}\sigma_0}(x, gx).$$

Since each $g^{-i}\sigma_0$ is the geodesic ray in a geodesic $g^{-i}[y_{g^{-1}}, y_g]$ between $y_{g^{-1}}$ and y_g , it follows from Lemma 6.10 that $|\beta_{g^{-i}\sigma_0}(x, gx) - \beta_{\sigma_0}(x, gx)| \leq 12\delta$. This implies

$$|\beta_{\sigma_0}(x, g^n x) - n\beta_{\sigma_0}(x, gx)| \leq 12\delta(n-1).$$

Hence we have

$$\left| \beta_{\sigma_0}(x, gx) - \frac{d_Z(x, g^n x)}{n} \right| \leq 12\delta.$$

Since this holds for all large $n \geq 1$, taking $n \rightarrow \infty$ yields the first claim.

Let us now show the last claim. We have

$$\beta_{\sigma}(w, gw) = \beta_{\sigma}(x, gx) + \beta_{\sigma}(w, x) - \beta_{g^{-1}\sigma}(w, x).$$

Since $g^{-1}\sigma(\infty) = y_g$ as well, it follows from Lemma 6.14 that

$$|\beta_{\sigma}(w, gw) - \beta_{\sigma}(x, gx)| \leq 40\delta.$$

Applying Lemma 6.14 again, we obtain

$$|\beta_{\sigma}(w, gw) - \beta_{\sigma_0}(x, gx)| \leq 80\delta.$$

By the first claim, we obtain

$$|\beta_{\sigma}(w, gw) - \ell(g)| \leq 92\delta$$

as desired. \square

If a subgroup $\Gamma < \text{Isom}(Z)$ acts properly discontinuously on Z , then the actions of Γ on ∂Z and \bar{Z} are convergence group actions [7, Lemma 2.11]. We denote by $\Lambda^Z = \Lambda_{\Gamma}^Z$ the limit set of Γ , which is the set of accumulation points of the Γ -orbit in ∂Z . We say that Γ is non-elementary if $\#\Lambda^Z \geq 3$.

7. ESSENTIAL SUBGROUPS FOR GRAPH-CONFORMAL MEASURES

Let G_1 be a connected semisimple real algebraic group and $\Gamma < G_1$ be a Zariski dense θ_1 -hypertransverse subgroup. Let (Z, d_Z) be a proper geodesic δ -hyperbolic space on which Γ acts properly discontinuously by isometries, with the Γ -equivariant homeomorphism $\iota : \Lambda^Z \rightarrow \Lambda^{\theta_1}$.

We keep the same notations as in Section 5. Let $\rho : \Gamma \rightarrow G_2$ be a Zariski dense θ_2 -regular faithful representation with a pair of ρ -equivariant continuous maps $f : \Lambda^{\theta_1} \rightarrow \Lambda_{\rho(\Gamma)}^{\theta_2}$ and $f_1 : \Lambda^{i(\theta_1)} \rightarrow \Lambda_{\rho(\Gamma)}^{i(\theta_2)}$. Let $G = G_1 \times G_2$ and consider the self-joining

$$\Gamma_\rho = (\text{id} \times \rho)(\Gamma) < G.$$

Its limit set in \mathcal{F}_θ is the graph $\Lambda_\rho^\theta = (\text{id} \times f)(\Lambda^{\theta_1})$. Recall that for a Γ -conformal measure ν on Λ^{θ_1} , the graph-conformal measure is defined as follows:

$$\nu_\rho = (\text{id} \times f)_* \nu.$$

The main goal of this section is to prove:

Theorem 7.1. *Let ν be a Γ -conformal measure of divergence type and ν_ρ the associated graph-conformal measure of Γ_ρ . If Γ_ρ is Zariski dense, then*

$$\mathbf{E}_{\nu_\rho}^\theta = \mathbf{a}_\theta.$$

Visual balls in Λ_ρ^θ . Via the equivariant homeomorphisms $\iota : \Lambda^Z \rightarrow \Lambda^{\theta_1}$ and $(\text{id} \times f) : \Lambda^{\theta_1} \rightarrow \Lambda_\rho^\theta$, we identify Λ^Z , Λ^{θ_1} and Λ_ρ^θ . In particular, we can use the notion of visual balls in Section 6 on all three spaces. More precisely, setting $f_0 := (\text{id} \times f) \circ \iota$, we can define the visual metric on Λ_ρ^θ as follows: for $x \in Z$ and $\xi, \eta \in \Lambda_\rho^\theta$,

$$d_x(\xi, \eta) := e^{-2\langle f_0^{-1}(\xi), f_0^{-1}(\eta) \rangle_x}.$$

We also use the same notation for the d_x -balls: for $\xi \in \Lambda_\rho^\theta$ and $r > 0$, $B_x(\xi, r) := \{\eta \in \Lambda_\rho^\theta : d_x(\xi, \eta) < r\}$. This allows us to regard Λ_ρ^θ as the limit set of Γ in ∂Z , and to employ the properties of visual metrics discussed in Section 6.

Main proposition. Since $(\text{id} \times \rho) : \Gamma \rightarrow \Gamma_\rho$ is an isomorphism, we can regard the Γ -action on Z as the Γ_ρ -action on Z : for $g \in \Gamma$ and $x \in Z$, $(g, \rho(g)) \cdot x = g \cdot x$. We will keep using this identification to ease the notations.

Proposition 7.2. *Let ν be a Γ -conformal measure of divergence type, and ν_ρ the associated graph-conformal measure of Γ_ρ . Let $\gamma_0 \in \Gamma_\rho$ be a loxodromic element such that $\ell(\gamma_0) > \frac{1}{2}(344\delta + 10^{100}\delta + \log 3)$. For any $\varepsilon > 0$ and a Borel subset $B \subset \mathcal{F}_\theta$ with $\nu_\rho(B) > 0$, there exists $\gamma \in \Gamma_\rho$ such that*

$$B \cap \gamma \gamma_0 \gamma^{-1} B \cap \left\{ \xi \in \Lambda_\rho^\theta : \|\beta_\xi^\theta(o, \gamma \gamma_0 \gamma^{-1} o) - \lambda_\theta(\gamma_0)\| < \varepsilon \right\}$$

has a positive ν_ρ -measure. In particular,

$$\lambda_\theta(\gamma_0) \in \mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho).$$

Remark 7.3. The proof of Proposition 7.2 is motivated by Robin [36] and Lee-Oh [32]. In both ([36], [32]) there exists a nice Busemann function due to the CAT(-1) and the higher rank Morse lemma respectively. In contrast, in the generality of our setting, the Busemann function on a Gromov hyperbolic space (Z, d_Z) is not as good as the one in CAT(-1) spaces, and the higher rank Morse lemma is not available. We overcome this difficulty by simultaneously controlling both the coarsely defined Busemann function on (Z, d_Z) and the \mathfrak{a}_θ -valued Busemann map on the higher rank symmetric space X to make the modified argument work. Our arguments are based on the dynamical properties of transverse subgroups and the uniformity and stability results on Busemann functions on (Z, d_Z) obtained in Section 6.

In the rest of this section, we fix a loxodromic element $\gamma_0 \in \Gamma_\rho$ and assume that $\ell(\gamma_0) > \frac{1}{2}(344\delta + 10^{100}\delta + \log 3)$. We denote by $\xi_0 \in \Lambda_\rho^\theta$ and $\eta \in \Lambda_\rho^{i(\theta)}$ the attracting and repelling fixed points of γ_0 respectively, which are identified with the attracting and repelling fixed points $y_{\gamma_0}, y_{\gamma_0^{-1}} \in \Lambda^Z$ respectively. Since ξ_0 and η are in general position, we can choose $p = go \in X$ where $g \in G$ is such that $\xi_0 = gP_\theta$ and $\eta = gw_0P_{i(\theta)}$. We also fix a point $x \in Z$ on a geodesic $[\xi_0, \eta]$ between ξ_0 and η in Z and a geodesic ray $\sigma_0 : [0, \infty) \rightarrow [\xi_0, \eta]$ with $\sigma_0(0) = x$ and $\sigma_0(\infty) = \xi_0$. Finally, we fix $0 < \varepsilon < 1/2$.

Covering the Myrberg limit set. We first make the choice of two constants:

$$C_1 = 10^{100}\delta \quad \text{and} \quad C_2 = 10^{10}\delta.$$

We then have

$$C_1 \geq 5C_2 + 104\delta \quad \text{and} \quad \ell(\gamma_0) > \frac{1}{2}(344\delta + C_1 + \log 3).$$

We only need these two properties; one can choose different C_1 and C_2 as long as they satisfy the above inequalities.

For each $\gamma \in \Gamma_\rho$, let $r_0(\gamma) > 0$ be the supremum of $r \geq 0$ such that

- we have

$$\sup_{\xi \in B_x(\gamma\xi_0, 3e^{8\delta}r)} \|\beta_\xi^\theta(p, \gamma\gamma_0^{\pm 1}\gamma^{-1}p) \mp \lambda_\theta(\gamma_0)\| < \varepsilon; \quad \text{and}$$

- for any geodesic ray $\sigma : [0, \infty) \rightarrow Z$ with $\sigma(\infty) \in B_x(\gamma\xi_0, 3e^{8\delta}r)$,

$$|\beta_\sigma(x, \gamma\gamma_0^{\pm 1}\gamma^{-1}x) \mp \ell(\gamma_0)| < C_1.$$

Such an r exists by Lemma 3.2, Lemma 6.20, and Lemma 6.15.

For each $R > 0$, we define

$$\mathcal{B}_R(\gamma_0, \varepsilon) = \{B_x(\gamma\xi_0, r) : \gamma \in \Gamma_\rho, 0 < r \leq \min(R, r_0(\gamma))\}.$$

Choose $0 < s = s(\gamma_0) < R$ small enough so that

- we have

$$\sup_{\xi \in B_x(\xi_0, s)} \|\beta_\xi^\theta(p, \gamma_0^{\pm 1}p) \mp \lambda_\theta(\gamma_0)\| < \frac{\varepsilon}{4};$$

- for any geodesic ray $\sigma : [0, \infty) \rightarrow Z$ with $\sigma(\infty) \in B_x(\xi_0, s)$,

$$|\beta_\sigma(x, \gamma_0^{\pm 1}x) \mp \ell(\gamma_0)| < C_2; \text{ and}$$

- we have

$$B_x(\xi_0, e^{2\ell(\gamma_0)+C_2}s) \subset O_{\varepsilon/(8\kappa)}^\theta(\eta, p) \cap O_{C_2}^Z(\eta, x)$$

where $\kappa > 0$ is the constant given in Lemma 3.3.

For each $\gamma \in \Gamma_\rho$ and $r > 0$, we set

$$D(\gamma\xi_0, r) = B_x\left(\gamma\xi_0, \frac{1}{3e^{8\delta}}e^{-2d_Z(x, \gamma x)}r\right).$$

Proposition 7.4. *Fix $R > 0$. Let $\xi \in \Lambda_\rho^\theta$ and $\gamma_i \in \Gamma_\rho$ be a sequence such that $\gamma_i^{-1}p \rightarrow \eta$ and $\gamma_i^{-1}\xi \rightarrow \xi_0$ as $i \rightarrow \infty$. Then for any $0 < r \leq e^{-500\delta}s(\gamma_0)$, there exists i_0 such that for all $i \geq i_0$, we have*

$$D(\gamma_i\xi_0, r) \in \mathcal{B}_R(\gamma_0, \varepsilon) \quad \text{and} \quad \xi \in D(\gamma_i\xi_0, r).$$

In particular, for any $R > 0$, we have

$$\Lambda_{\rho, M}^\theta \subset \bigcup_{D \in \mathcal{B}_R(\gamma_0, \varepsilon)} D.$$

Proof. We first claim that $D(\gamma_i\xi_0, r) \in \mathcal{B}_R(\gamma_0, \varepsilon)$ for all large i . By Lemma 2.5 and the equivariance of the homeomorphism $\Lambda^Z \rightarrow \Lambda^{i(\theta_1)}$, we have that $\gamma_i^{-1}x \rightarrow y_{\gamma_0^{-1}}$ in \bar{Z} , noting that $\eta = y_{\gamma_0^{-1}}^{i(\theta)}$. Hence $O_{C_2}^Z(y_{\gamma_0^{-1}}, x) \subset O_{C_2+2\delta}^Z(\gamma_i^{-1}x, x)$ for all i by Lemma 6.19.

For each $i \geq 1$, we set $s_i = \frac{1}{3e^{8\delta}}e^{-2d_Z(\gamma_i x, x)}r$. We need to show that

$$\sup_{\xi' \in B_x(\gamma_i\xi_0, 3e^{8\delta}s_i)} \|\beta_{\xi'}^\theta(p, \gamma_i\gamma_0^{\pm 1}\gamma_i^{-1}p) \mp \lambda_\theta(\gamma_0)\| < \varepsilon$$

and for any geodesic ray $\sigma : [0, \infty) \rightarrow Z$ with $\sigma(\infty) \in B_x(\gamma_i\xi_0, 3e^{8\delta}s_i)$,

$$|\beta_\sigma(x, \gamma_i\gamma_0^{\pm 1}\gamma_i^{-1}x) \mp \ell(\gamma_0)| < C_1.$$

Let $\xi' \in B_x(\gamma_i\xi_0, 3e^{8\delta}s_i)$ and $\sigma : [0, \infty) \rightarrow Z$ a geodesic ray with $\sigma(\infty) = \xi'$. We have from Lemma 6.17 that

$$\begin{aligned} d_x(\xi_0, \gamma_i^{-1}\xi') &= d_{\gamma_i x}(\gamma_i\xi_0, \xi') \leq e^{244\delta}e^{\beta_{\gamma_i\sigma_0}(x, \gamma_i x) + \beta_\sigma(x, \gamma_i x)}d_x(\gamma_i\xi_0, \xi') \\ &< e^{244\delta}e^{2d_Z(x, \gamma_i x)}e^{-2d_Z(\gamma_i x, x)}r = e^{244\delta}r. \end{aligned}$$

Since $e^{244\delta}r \leq s(\gamma_0)$, we have

$$\|\beta_{\gamma_i^{-1}\xi'}^\theta(p, \gamma_0 p) - \lambda_\theta(\gamma_0)\| < \frac{\varepsilon}{4} \quad \text{and} \quad |\beta_{\gamma_i^{-1}\sigma}(x, \gamma_0 x) - \ell(\gamma_0)| < C_2.$$

Hence we have

$$\begin{aligned} d_x(\xi_0, \gamma_0^{-1}\gamma_i^{-1}\xi') &= d_{\gamma_0 x}(\xi_0, \gamma_i^{-1}\xi') \leq e^{244\delta}e^{\beta_{\sigma_0}(x, \gamma_0 x) + \beta_{\gamma_i^{-1}\sigma}(x, \gamma_0 x)}d_x(\xi_0, \gamma_i^{-1}\xi') \\ &\leq e^{244\delta}e^{12\delta}e^{2\ell(\gamma_0)+C_2}e^{244\delta}r \end{aligned}$$

by Lemma 6.20. Since $e^{500\delta}r < s(\gamma_0)$, this implies that $\gamma_i^{-1}\xi', \gamma_0^{-1}\gamma_i^{-1}\xi' \in B_x(\xi_0, e^{2\ell(\gamma_0)+C_2}s)$. Since $B_x(\xi_0, e^{2\ell(\gamma_0)+C_2}s) \subset O_{\varepsilon/(8\kappa)}^\theta(\eta, p)$ and $\gamma_i^{-1}p \rightarrow \eta$, we obtain from Corollary 3.4 that

$$\|\beta_{\gamma_i^{-1}\xi'}^\theta(\gamma_i^{-1}p, p) - \beta_{\gamma_0^{-1}\gamma_i^{-1}\xi'}^\theta(\gamma_i^{-1}p, p)\| < \frac{\varepsilon}{2} \quad \text{for all but finitely many } i.$$

Now we have

$$\begin{aligned} & \|\beta_{\xi'}^\theta(p, \gamma_i\gamma_0\gamma_i^{-1}p) - \lambda_\theta(\gamma_0)\| \\ &= \|\beta_{\xi'}^\theta(p, \gamma_i p) + \beta_{\xi'}^\theta(\gamma_i p, \gamma_i\gamma_0 p) + \beta_{\xi'}^\theta(\gamma_i\gamma_0 p, \gamma_i\gamma_0\gamma_i^{-1}p) - \lambda_\theta(\gamma_0)\| \\ &\leq \|\beta_{\xi'}^\theta(p, \gamma_i p) - \beta_{\xi'}^\theta(\gamma_i\gamma_0\gamma_i^{-1}p, \gamma_i\gamma_0 p)\| + \|\beta_{\xi'}^\theta(\gamma_i p, \gamma_i\gamma_0 p) - \lambda_\theta(\gamma_0)\| \\ &= \|\beta_{\gamma_i^{-1}\xi'}^\theta(\gamma_i^{-1}p, p) - \beta_{\gamma_0^{-1}\gamma_i^{-1}\xi'}^\theta(\gamma_i^{-1}p, p)\| + \|\beta_{\gamma_i^{-1}\xi'}^\theta(p, \gamma_0 p) - \lambda_\theta(\gamma_0)\| \\ &< \varepsilon/2 + \varepsilon/4 < \varepsilon. \end{aligned}$$

Similarly, as $B_x(\xi_0, e^{2\ell(\gamma_0)+C_2}s) \subset O_{C_2}^Z(y_{\gamma_0^{-1}}, x)$, it follows from Lemma 6.19 and Lemma 6.18 that

$$|\beta_{\gamma_i^{-1}\sigma}(\gamma_i^{-1}x, x) - \beta_{\gamma_0^{-1}\gamma_i^{-1}\sigma}(\gamma_i^{-1}x, x)| < 4C_2 + 104\delta \quad \text{for all large } i.$$

Hence we have

$$\begin{aligned} & |\beta_\sigma(x, \gamma_i\gamma_0\gamma_i^{-1}x) - \ell(\gamma_0)| \\ &\leq |\beta_{\gamma_i^{-1}\sigma}(\gamma_i^{-1}x, x) - \beta_{\gamma_0^{-1}\gamma_i^{-1}\sigma}(\gamma_i^{-1}x, x)| + |\beta_{\gamma_i^{-1}\sigma}(x, \gamma_0x) - \ell(\gamma_0)| \\ &< 4C_2 + 104\delta + C_2 \leq C_1. \end{aligned}$$

By the same argument, we also have

$$\|\beta_{\xi'}^\theta(p, \gamma_i\gamma_0^{-1}\gamma_i^{-1}p) + \lambda_\theta(\gamma_0)\| < \varepsilon \quad \text{and} \quad |\beta_\sigma(x, \gamma_i\gamma_0^{-1}\gamma_i^{-1}x) + \ell(\gamma_0)| < C_1.$$

Since $\xi' \in B_x(\gamma_i\xi_0, 3e^{8\delta}s_i)$ and the geodesic ray σ were arbitrary, it shows $D(\gamma_i\xi_0, r) \in \mathcal{B}_R(\gamma_0, \varepsilon)$ for all large i .

We now prove the second claim that $\xi \in D(\gamma_i\xi_0, r)$ for all large i . Since $\gamma_i^{-1}\xi \rightarrow \xi_0$, we may assume that

$$\gamma_i^{-1}\xi \in B_x(\xi_0, e^{2\ell(\gamma_0)+C_2}s) \subset O_{C_2}^Z(y_{\gamma_0^{-1}}, x) \quad \text{for all } i \geq 1.$$

By Lemma 6.19, we have

$$\gamma_i^{-1}\xi \in O_{C_2+2\delta}^Z(\gamma_i^{-1}x, x) \quad \text{for all large } i.$$

Note that $\xi_0 \in O_{C_2+2\delta}^Z(\gamma_i^{-1}x, x)$ as well. Let σ be a geodesic ray with $\sigma(\infty) = \xi$. It follows from Lemma 6.18 that

$$\begin{aligned} |\beta_{\gamma_i^{-1}\sigma}(\gamma_i^{-1}x, x) - d_Z(\gamma_i^{-1}x, x)| &< 2C_2 + 52\delta; \\ |\beta_{\sigma_0}(\gamma_i^{-1}x, x) - d_Z(\gamma_i^{-1}x, x)| &< 2C_2 + 52\delta. \end{aligned}$$

Therefore, we have

$$\begin{aligned} d_x(\gamma_i \xi_0, \xi) &= d_{\gamma_i^{-1}x}(\xi_0, \gamma_i^{-1}\xi) \\ &\leq e^{244\delta} e^{-(\beta_{\sigma_0}(\gamma_i^{-1}x, x) + \beta_{\gamma_i^{-1}\sigma}(\gamma_i^{-1}x, x))} d_x(\xi_0, \gamma_i^{-1}\xi) \\ &< e^{244\delta} e^{-2d_Z(\gamma_i^{-1}x, x)} e^{4C_2 + 104\delta} d_x(\xi_0, \gamma_i^{-1}\xi). \end{aligned}$$

Since $\gamma_i^{-1}\xi \rightarrow \xi_0$, we have $d_x(\xi_0, \gamma_i^{-1}\xi) < e^{-(4C_2 + 348\delta)} \frac{1}{3e^{8\delta}} r$ for all large i , and hence $\xi \in D(\gamma_i \xi_0, r)$, completing the proof. \square

Approximation by d_x -balls. From now on, let ν be a (Γ, ψ) -conformal measure of divergence type and ν_ρ the associated graph-conformal measure of Γ_ρ . Note that ν_ρ is a $(\Gamma_\rho, \sigma_\psi)$ -conformal measure where $\sigma_\psi \in \mathfrak{a}_\theta^*$ is the composition $\psi \circ p_{\theta_1}$ (Proposition 4.4). It is more convenient to use the following conformal measure ν_p (with respect to the basepoint p) as $\mathbb{E}_{\nu_p}^\theta(\Gamma_\rho) = \mathbb{E}_{\nu_\rho}^\theta(\Gamma_\rho)$:

$$d\nu_p(\xi) = e^{\sigma_\psi(\beta_\xi^{(\circ, p)})} d\nu_\rho(\xi).$$

Proposition 7.5. *Let $B \subset \mathcal{F}_\theta$ be a Borel subset with $\nu_p(B) > 0$. Then for ν_p -a.e. $\xi \in B$, we have*

$$\lim_{R \rightarrow 0} \sup_{\xi \in D, D \in \mathcal{B}_R(\gamma_0, \varepsilon)} \frac{\nu_p(B \cap D)}{\nu_p(D)} = 1.$$

Proof. For a Borel function $h : \mathcal{F}_\theta \rightarrow \mathbb{R}$, define $h^* : \mathcal{F}_\theta \rightarrow \mathbb{R}$ as

$$h^*(\xi) := \lim_{R \rightarrow 0} \sup_{\xi \in D, D \in \mathcal{B}_R(\gamma_0, \varepsilon)} \frac{1}{\nu_p(D)} \int_D h d\nu_p.$$

By Proposition 7.4, h^* is well-defined on $\Lambda_{\rho, M}^\theta$. Since $\Lambda_{\rho, M}^\theta$ has full ν_p -measure by Theorem 5.3, h^* is well-defined for ν_p -a.e. $\xi \in \mathcal{F}_\theta$. It suffices to show that $h(\xi) = h^*(\xi)$ for ν_p -a.e. $\xi \in \mathcal{F}_\theta$; taking $h = \mathbb{1}_B$ implies the desired identity. Note that $h = h^*$ if h is continuous; we now consider the general case.

Claim. We claim that for any $\alpha > 0$, we have

$$\nu_p(h^* > \alpha) \leq \frac{e^{\sigma_\psi(\lambda_\theta(\gamma_0)) + \|\sigma_\psi\|\varepsilon}}{\alpha} \int_{\mathcal{F}_\theta} |h| d\nu_p.$$

To see this, it suffices to show that for any compact $Q \subset \{h^* > \alpha\}$, we have

$$\nu_p(Q) \leq \frac{e^{\sigma_\psi(\lambda_\theta(\gamma_0)) + \|\sigma_\psi\|\varepsilon}}{\alpha} \int_{\mathcal{F}_\theta} |h| d\nu_p.$$

Fix $R > 0$ and a compact subset $Q \subset \{h^* > \alpha\}$. By the definition of h^* , for each $q \in Q$, there exists $D_q \in \mathcal{B}_R(\gamma_0, \varepsilon)$ containing q such that

$$\frac{1}{\nu_p(D_q)} \int_{D_q} h d\nu_p > \alpha.$$

Since Q is compact, we have a finite subcover $\{D_i = B_x(\gamma_i \xi_0, s_i)\}$ of $\{D_q : q \in Q\}$, where $\gamma_i \in \Gamma_\rho$ and $s_i = \frac{1}{3e^{8\delta}} e^{-2d_Z(\gamma_i^{-1}x, x)} r_i$ for some $r_i > 0$.

By Lemma 6.13, there exists a subcollection D_{i_1}, \dots, D_{i_k} of disjoint subsets such that

$$\bigcup_i D_i \subset \bigcup_{j=1}^k 3e^{8\delta} D_{i_j}$$

where $3e^{8\delta} D_{i_j} = B_x(\gamma_{i_j} \xi_0, 3e^{8\delta} s_{i_j})$.

We observe that for each j , $3e^{8\delta} D_{i_j} \subset \gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} D_{i_j}$. Indeed, for $\xi \in 3e^{8\delta} D_{i_j}$ and a geodesic ray σ with $\sigma(\infty) = \xi$, we have from Lemma 6.17, $\xi \in 3e^{8\delta} D_{i_j}$, and $D_{i_j} \in \mathcal{B}_R(\gamma_0, \varepsilon)$ that

$$\begin{aligned} d_x(\gamma_{i_j} \xi_0, \gamma_{i_j} \gamma_0 \gamma_{i_j}^{-1} \xi) &= d_{\gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} x}(\gamma_{i_j} \xi_0, \xi) \\ &\leq e^{244\delta} e^{-(\beta_{\gamma_{i_j} \sigma_0}(\gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} x, x) + \beta_\sigma(\gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} x, x))} d_x(\gamma_{i_j} \xi_0, \xi) \\ &\leq e^{244\delta} e^{92\delta} e^{-2\ell(\gamma_0) + C_1} 3e^{8\delta} s_{i_j} < s_{i_j}. \end{aligned}$$

This shows $\gamma_{i_j} \gamma_0 \gamma_{i_j}^{-1} \xi \in D_{i_j}$, and hence $\xi \in \gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} D_{i_j}$.

Therefore, we have

$$\begin{aligned} \nu_p(3e^{8\delta} D_{i_j}) &\leq \nu_p(\gamma_{i_j} \gamma_0^{-1} \gamma_{i_j}^{-1} D_{i_j}) = \int_{D_{i_j}} e^{\sigma_\psi(\beta_\xi^\theta(p, \gamma_{i_j} \gamma_0 \gamma_{i_j}^{-1} p))} d\nu_p(\xi) \\ &\leq e^{\sigma_\psi(\lambda_\theta(\gamma_0)) + \|\sigma_\psi\| \varepsilon} \nu_p(D_{i_j}). \end{aligned}$$

Now it follows that

$$\begin{aligned} \nu_p(Q) &\leq \sum_{j=1}^k \nu_p(3e^{8\delta} D_{i_j}) \\ &\leq \sum_{j=1}^k \frac{e^{\sigma_\psi(\lambda_\theta(\gamma_0)) + \|\sigma_\psi\| \varepsilon}}{\alpha} \int_{D_{i_j}} h d\nu_p \leq \frac{e^{\sigma_\psi(\lambda_\theta(\gamma_0)) + \|\sigma_\psi\| \varepsilon}}{\alpha} \int_{\mathcal{F}_\theta} |h| d\nu_p, \end{aligned}$$

as desired.

We now finish the proof of the proposition by showing that $h(\xi) = h^*(\xi)$ for ν_p -a.e. ξ . We first show that $h(\xi) \leq h^*(\xi)$ for ν_p -a.e. ξ . Let $\alpha > 0$ and a sequence of continuous functions $h_n \rightarrow h$ in $L^1(\nu_p)$. Since h_n is continuous, $h_n^* = h_n$. Now we have

$$\begin{aligned} \nu_p(h - h^* > \alpha) &\leq \nu_p(h - h_n > \alpha/2) + \nu_p(h_n^* - h^* > \alpha/2) \\ &\leq \frac{2}{\alpha} \|h - h_n\|_{L^1} + \frac{2}{\alpha} e^{\sigma_\psi(\lambda_\theta(\gamma_0)) + \|\sigma_\psi\| \varepsilon} \|h - h_n\|_{L^1}. \end{aligned}$$

Since $\|h - h_n\|_{L^1} \rightarrow 0$ as $n \rightarrow \infty$, we have $\nu_p(h - h^* > \alpha) = 0$. Since $\alpha > 0$ is arbitrary, $h(\xi) \leq h^*(\xi)$ for ν_p -a.e. ξ . The similar argument shows $h^*(\xi) \leq h(\xi)$ for ν_p -a.e. ξ , and it completes the proof. \square

Proof of Proposition 7.2. Let $B \subset \mathcal{F}_\theta$ be a Borel subset with $\nu_p(B) > 0$. It suffices to show that for some $\gamma \in \Gamma_\rho$, the set

$$B \cap \gamma\gamma_0\gamma^{-1}B \cap \{\xi : \|\beta_\xi^\theta(p, \gamma\gamma_0\gamma^{-1}p) - \lambda_\theta(\gamma_0)\| < \varepsilon\}$$

has positive ν_p -measure.

By Proposition 7.5, there exists $D = B_x(\gamma\xi_0, r) \in \mathcal{B}_R(\gamma_0, \varepsilon)$ such that

$$(7.1) \quad \nu_p(D \cap B) > (1 + e^{-\sigma_\psi(\lambda_\theta(\gamma_0)) - \|\sigma_\psi\|\varepsilon})^{-1} \nu_p(D).$$

Since $r < r_0(\gamma)$, we have

$$D \subset \{\xi : \|\beta_\xi^\theta(p, \gamma\gamma_0^\pm\gamma^{-1}p) \mp \lambda_\theta(\gamma_0)\| < \varepsilon\}$$

and for any geodesic ray σ with $\sigma(\infty) \in D$, we have

$$|\beta_\sigma(x, \gamma\gamma_0^\pm\gamma^{-1}x) \mp \ell(\gamma_0)| < C_1.$$

This implies

$$B \cap \gamma\gamma_0\gamma^{-1}B \cap \{\xi : \|\beta_\xi^\theta(p, \gamma\gamma_0\gamma^{-1}p) - \lambda_\theta(\gamma_0)\| < \varepsilon\} \supset (D \cap B) \cap \gamma\gamma_0\gamma^{-1}(D \cap B).$$

Hence it suffices to show

$$(7.2) \quad \nu_p((D \cap B) \cap \gamma\gamma_0\gamma^{-1}(D \cap B)) > 0.$$

By the conformality, we have

$$\begin{aligned} \nu_p(\gamma\gamma_0\gamma^{-1}(D \cap B)) &= \int_{D \cap B} e^{\sigma_\psi(\beta_\xi^\theta(p, \gamma\gamma_0^{-1}\gamma^{-1}p))} d\nu_p(\xi) \\ &> e^{-\sigma_\psi(\lambda_\theta(\gamma_0)) - \|\sigma_\psi\|\varepsilon} \nu_p(D \cap B). \end{aligned}$$

Hence we have

$$\nu_p(D \cap B) + \nu_p(\gamma\gamma_0\gamma^{-1}(D \cap B)) > (1 + e^{-\sigma_\psi(\lambda_\theta(\gamma_0)) - \|\sigma_\psi\|\varepsilon}) \nu_p(D \cap B).$$

Together with the choice (7.1) of D , we obtain

$$(7.3) \quad \nu_p(D \cap B) + \nu_p(\gamma\gamma_0\gamma^{-1}(D \cap B)) > \nu_p(D).$$

We claim that $\gamma\gamma_0\gamma^{-1}D \subset D$. Indeed, if $\xi \in D$ and σ is a geodesic ray with $\sigma(\infty) = \xi$, then by Lemma 6.17 and Lemma 6.20,

$$\begin{aligned} d_x(\gamma\xi_0, \gamma\gamma_0\gamma^{-1}\xi) &= d_{\gamma\gamma_0^{-1}\gamma^{-1}x}(\gamma\xi_0, \xi) \\ &\leq e^{244\delta} e^{\beta_{\gamma\sigma_0}(x, \gamma\gamma_0^{-1}\gamma^{-1}x) + \beta_\sigma(x, \gamma\gamma_0^{-1}\gamma^{-1}x)} d_x(\gamma\xi_0, \xi) \\ &< e^{244\delta} e^{92\delta} e^{-2\ell(\gamma_0) + C_1 r} < r. \end{aligned}$$

Hence the claim follows.

Now both $D \cap B$ and $\gamma\gamma_0\gamma^{-1}(D \cap B)$ are subsets of D . Therefore, (7.3) must imply (7.2), completing the proof of Proposition 7.2. \square

Corollary 7.6. *For any loxodromic $\gamma_0 \in \Gamma_\rho$, we have*

$$\lambda_\theta(\gamma_0) \in \mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho).$$

Proof. Let $\gamma_0 \in \Gamma$ be a loxodromic element. For sufficiently large n , both γ_0^n and γ_0^{n+1} satisfy the condition of Proposition 7.2. Hence we have

$$n\lambda_\theta(\gamma_0), (n+1)\lambda_\theta(\gamma_0) \in \mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho).$$

Since $\mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho)$ is a subgroup of \mathfrak{a}_θ , we have

$$\lambda_\theta(\gamma_0) = (n+1)\lambda_\theta(\gamma_0) - n\lambda_\theta(\gamma_0) \in \mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho).$$

□

Proof of Theorem 7.1. By Corollary 7.6, $\lambda_\theta(\gamma_0) \in \mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho)$ for all loxodromic $\gamma_0 \in \Gamma_\rho$. Since $\mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho)$ is a closed subgroup of \mathfrak{a}_θ , it follows from Theorem 2.3 that $\mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho) = \mathfrak{a}_\theta$ if Γ_ρ is Zariski dense. □

Essential subgroups for hypertransverse subgroups. The same argument applies to a Zariski dense θ -hypertransverse subgroup $\Gamma < G$, which is not necessarily a self-joining. Therefore we deduce:

Theorem 7.7. *Let G be a semisimple real algebraic group and $\Gamma < G$ a Zariski dense θ -hypertransverse subgroup. For a Γ -conformal measure ν of divergence type, we have*

$$\mathbf{E}_\nu^\theta(\Gamma) = \mathfrak{a}_\theta.$$

8. SINGULARITY OF THE GRAPH-CONFORMAL MEASURE

We are finally ready to prove our main rigidity theorems. We recall the setting: let G_1 and G_2 be connected simple real algebraic groups and $\Gamma < G_1$ be a Zariski dense θ_1 -hypertransverse subgroup with the limit set $\Lambda^{\theta_1} \subset \mathcal{F}_{\theta_1}$. Let $\rho : \Gamma \rightarrow G_2$ be a Zariski dense θ_2 -regular representation with ρ -equivariant continuous maps $f : \Lambda^{\theta_1} \rightarrow \mathcal{F}_{\theta_2}$ and $f_i : \Lambda^{i(\theta_1)} \rightarrow \mathcal{F}_{i(\theta_2)}$. Let ν be a (Γ, ψ) -conformal measure of divergence type, for $\psi \in \mathfrak{a}_{\theta_1}^*$.

Recall that $\Gamma_\rho = (\text{id} \times \rho)(\Gamma)$ is the self-joining of Γ via ρ which is a discrete subgroup of $G = G_1 \times G_2$. The graph-conformal measure $\nu_\rho = (\text{id} \times f)_* \nu$ is the unique $(\Gamma_\rho, \sigma_\psi)$ -conformal measure on Λ_ρ^θ where σ_ψ is the composition of ψ with the projection $\mathfrak{a}_\theta \rightarrow \mathfrak{a}_{\theta_1}$ (Proposition 4.4).

Theorem 8.1. *If Γ_ρ is Zariski dense, then*

$$\nu_\phi \not\ll \nu_\rho$$

for all (Γ_ρ, ϕ) -conformal measure ν_ϕ on \mathcal{F}_θ with $\phi \neq \sigma_\psi$.

Proof. Let ν_ϕ be a (Γ_ρ, ϕ) -conformal measure on \mathcal{F}_θ for some $\phi \in \mathfrak{a}_\theta^*$. Suppose that $\nu_\phi \ll \nu_\rho$. By Theorem 7.1, we have $\mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho) = \mathfrak{a}_\theta$. Hence it follows from Proposition 3.8 that $\phi = \sigma_\psi$ on $\mathbf{E}_{\nu_\rho}^\theta(\Gamma_\rho) = \mathfrak{a}_\theta$. Therefore, $\phi = \sigma_\psi$ and the theorem follows. □

Proof of Theorem 1.4. Suppose that Γ_ρ is Zariski dense and there exists a $(\rho(\Gamma), \varphi)$ -conformal measure ν_φ on $\Lambda_{\rho(\Gamma)}^{\theta_2}$ for some $\varphi \in \mathfrak{a}_{\theta_2}^*$ such that

$$\nu_\varphi \ll f_*\nu.$$

Then by Proposition 4.4, we have

$$(f^{-1} \times \text{id})_*\nu_\varphi \ll \nu_\rho$$

and $(f^{-1} \times \text{id})_*\nu_\varphi$ is a $(\Gamma_\rho, \sigma_\varphi)$ -conformal measure where σ_φ is the composition of the projection $\mathfrak{a}_\theta \rightarrow \mathfrak{a}_{\theta_2}$ and $\varphi \in \mathfrak{a}_{\theta_2}^*$. By Theorem 8.1, we must have $\sigma_\psi = \sigma_\varphi$.

On the other hand, $\mathfrak{a}_{\theta_1} < \mathfrak{a}_\theta = \mathfrak{a}_{\theta_1} \oplus \mathfrak{a}_{\theta_2}$ is contained in $\ker \sigma_\varphi$ while $\sigma_\psi(u) = \psi(u) \neq 0$ for some $u \in \mathfrak{a}_{\theta_1}$, which is a contradiction. Therefore, Γ_ρ is not Zariski dense and hence ρ extends to a Lie group isomorphism $G_1 \rightarrow G_2$ by Lemma 4.2. \square

Proof of Theorem 1.6. By Theorem 7.7, we have $\mathbf{E}_\nu^\theta(\Gamma) = \mathfrak{a}_\theta$. Hence Theorem 1.6 follows by the same argument as in the proof of Theorem 8.1. \square

9. DEFORMATIONS OF TRANSVERSE REPRESENTATIONS

In this section, we consider deformations of transverse representations to which Theorem 8.1 can be applied. We keep the same notations from previous sections. Let (Z, d_Z) be a proper geodesic δ -hyperbolic space and $\Delta < \text{Isom}(Z)$ a non-elementary subgroup acting properly discontinuously on Z . For $i = 1, 2$, we consider θ_i -transverse representations $\rho_i : \Delta \rightarrow G_i$ and write $\Gamma_i := \rho_i(\Delta)$. The conjugate $\rho = \rho_2 \circ \rho_1|_\Delta^{-1}$ between two representations is referred to as a deformation from ρ_1 to ρ_2 :

$$\begin{array}{ccc} & & \Gamma_1 \\ & \nearrow^{\rho_1} & \downarrow \rho \\ \Delta & & \Gamma_2 \\ & \searrow_{\rho_2} & \end{array}$$

In this setting, we obtain the following stronger form of the conformal measure rigidity theorem which was stated as Theorem 1.8 in the introduction:

Theorem 9.1. *There exists a pair of ρ -boundary maps $f : \Lambda_{\Gamma_1}^{\theta_1} \rightarrow \mathcal{F}_{\theta_2}$ and $f_i : \Lambda_{\Gamma_1}^{i(\theta_1)} \rightarrow \mathcal{F}_{i(\theta_2)}$. Moreover, unless $\rho : \Gamma_1 \rightarrow \Gamma_2$ does not extend to a Lie group isomorphism $G_1 \rightarrow G_2$,*

$$\nu_2 \ll f_*\nu_1$$

for any Γ_1 -conformal measure ν_1 of divergence type and Γ_2 -conformal measure ν_2 . In particular, if ν_2 is further assumed to be of divergence type, then

$$\nu_2 \perp f_*\nu_1.$$

Proof. By definition of the transverse representation, for $i = 1, 2$, we have a ρ_i -equivariant homeomorphism $f_i : \Lambda_\Delta^Z \rightarrow \Lambda_{\Gamma_i}^{\theta_i \cup i(\theta_i)}$. Together with the canonical projections $\Lambda_{\Gamma_i}^{\theta_i \cup i(\theta_i)} \rightarrow \Lambda_{\Gamma_i}^{\theta_i}$ and $\Lambda_{\Gamma_i}^{\theta_i \cup i(\theta_i)} \rightarrow \Lambda_{\Gamma_i}^{i(\theta_i)}$, we have the following commutative diagram:

$$\begin{array}{ccccc}
 \Lambda_{\Gamma_1}^{i(\theta_1)} & \xleftarrow{\sim} & \Lambda_{\Gamma_1}^{\theta_1 \cup i(\theta_1)} & \xrightarrow{\sim} & \Lambda_{\Gamma_1}^{\theta_1} \\
 \downarrow f_i & & \uparrow f_1 & & \downarrow f \\
 & & \Lambda_\Delta^Z & & \\
 & & \downarrow f_2 & & \\
 \Lambda_{\Gamma_2}^{i(\theta_1)} & \xleftarrow{\sim} & \Lambda_{\Gamma_2}^{\theta_1 \cup i(\theta_1)} & \xrightarrow{\sim} & \Lambda_{\Gamma_2}^{\theta_1}
 \end{array}$$

As indicated in the above diagram, the projections $\Lambda_{\Gamma_i}^{\theta_i \cup i(\theta_i)} \rightarrow \Lambda_{\Gamma_i}^{\theta_i}$ and $\Lambda_{\Gamma_i}^{\theta_i \cup i(\theta_i)} \rightarrow \Lambda_{\Gamma_i}^{i(\theta_i)}$, $i = 1, 2$, are homeomorphisms due to the θ_i -antipodality of Γ_i [26, Lemma 9.5]. Hence the maps f and f_i are well-defined as above, and are homeomorphisms. Moreover, since all maps in the diagram are equivariant under the actions of the corresponding groups, f and f_i are ρ -equivariant. Therefore, they form a pair of ρ -boundary maps.

This allows us to apply Theorem 1.4, finishing the proof. \square

10. HOROSPHERICAL FOLIATIONS AND BURGER-ROBLIN MEASURES

In the rest of the paper, let G be a connected semisimple real algebraic group and fix a non-empty $\theta \subset \Pi$. In this section, we discuss ergodic properties of Burger-Roblin measures on horospherical foliations.

Recall the space

$$\mathcal{H}_\theta := \mathcal{F}_\theta \times \mathfrak{a}_\theta$$

and the actions of G and A_θ on \mathcal{H}_θ given as follows: for $(\xi, u) \in \mathcal{H}_\theta$, $g \in G$ and $a \in A_\theta$,

$$\begin{aligned}
 (10.1) \quad g \cdot (\xi, u) &= (g\xi, u + \beta_\xi^\theta(g^{-1}, e)); \\
 (\xi, u) \cdot a &= (\xi, u + \log a).
 \end{aligned}$$

Denoting by $g^+ = gP_\theta \in F_\theta$, the map $g \mapsto (g^+, \beta_{g^+}^\theta(e, g))$ induces a homeomorphism

$$G/N_\theta S_\theta \simeq \mathcal{H}_\theta.$$

Hence the space \mathcal{H}_θ can be considered as the θ -horospherical foliation. Indeed, when G is of rank one, \mathcal{H}_θ is the horospherical foliation of the unit tangent bundle of G/K .

Since A_θ normalizes $N_\theta S_\theta$, the quotient $G/N_\theta S_\theta$ admits both left G -action and right A_θ -action, and the above homeomorphism is (G, A_θ) -equivariant.

A Radon measure m on \mathcal{H}_θ is A_θ -semi-invariant if there exists a linear form $\chi_m \in \mathfrak{a}_\theta^*$ such that for all $a \in A_\theta$, we have

$$a_*m = e^{\chi_m(\log a)}m.$$

We define a Γ -invariant A_θ -semi-invariant Radon measure on \mathcal{H}_θ , called Burger-Roblin measure.

Definition 10.1 (Burger-Roblin measures). Let $\Gamma < G$ be a discrete subgroup and ν a (Γ, ψ) -conformal measure on \mathcal{F}_θ for some $\psi \in \mathfrak{a}_\theta^*$. The *Burger-Roblin measure* m_ν^{BR} on \mathcal{H}_θ associated to ν is defined by

$$dm_\nu^{\text{BR}}(\xi, u) := e^{\psi(u)}d\nu(\xi)du$$

where du is the Lebesgue measure on \mathfrak{a}_θ .

In fact, all Γ -invariant A_θ -semi-invariant measures arise as Burger-Roblin measures. See ([1], [10], [30]) for rank one settings, and [32, Proposition 10.25] for higher rank:

Proposition 10.2. *Let $\Gamma < G$ be a Zariski dense discrete subgroup. Any Γ -invariant A_θ -semi-invariant Radon measure on \mathcal{H}_θ is proportional a Burger-Roblin measure associated with some Γ -conformal measure on \mathcal{F}_θ .*

Ergodicity of horospherical foliations. We now prove the ergodicity of horospherical foliations with respect to Burger-Roblin measures. The size of the essential subgroup plays a role of criterion for the ergodicity of actions of horospherical foliations. The following was proved in [38] for abstract measurable dynamical systems, and more direct proof for particular case of CAT(−1) spaces was given in [36, Proposition 2.1]. Following [36], the higher rank version was obtained in [32, Proposition 9.2] when $\theta = \Pi$. The same proofs as in ([32] and [36]) works for general θ :

Proposition 10.3. *Let $\Gamma < G$ be a Zariski dense discrete subgroup and ν a Γ -conformal measure on \mathcal{F}_θ . The Γ -action on $(\mathcal{H}_\theta, m_\nu^{\text{BR}})$ is ergodic if and only if the Γ -action on $(\mathcal{F}_\theta, \nu)$ is ergodic and $E_\nu^\theta(\Gamma) = \mathfrak{a}_\theta$.*

Proof of Theorem 1.11. Let Γ be a Zariski dense θ -hypertransverse subgroup. Let ν be a Γ -conformal measure of divergence type. By Theorem 7.7, we have $E_\nu^\theta(\Gamma) = \mathfrak{a}_\theta$. Moreover, $(\mathcal{F}_\theta, \Gamma, \nu)$ is ergodic by Theorem 5.5. Therefore, the ergodicity of the Γ -action on $(\mathcal{H}_\theta, m_\nu^{\text{BR}})$ follows from Proposition 10.3. \square

Ergodic decomposition. In the rest of the section, we consider the case $\theta = \Pi$; we omit the subscripts and superscripts for $\theta = \Pi$. Let $\Gamma < G$ be a Zariski dense Π -hypertransverse subgroup.

For a (Γ, ψ) -conformal measure ν on \mathcal{F} for some $\psi \in \mathfrak{a}^*$, the associated Burger-Roblin measure \hat{m}_ν^{BR} on $\Gamma \backslash G$ is defined in (1.1). Let ν_i be a $(\Gamma, \psi \circ i)$ -conformal measure on \mathcal{F} . We now define the Bowen-Margulis-Sullivan

measure for the pair (ν, ν_1) . The generalized Hopf-parametrization for G is an isomorphism $G/M \rightarrow \mathcal{F}^{(2)} \times \mathfrak{a}$ defined by

$$gM \mapsto (g^+, g^-, \beta_{g^+}(e, g))$$

where $g^+ = gP$, $g^- = gw_0P \in \mathcal{F}$. By fixing a Borel section $G/M \rightarrow G$, it induces an isomorphism

$$(10.2) \quad G \rightarrow \mathcal{F}^{(2)} \times \mathfrak{a} \times M.$$

Via (10.2), the following defines a left Γ -invariant and right AM -invariant measure on G : for $g \in G$,

$$d\hat{m}_{\nu, \nu_1}^{\text{BMS}}(g) := e^{\psi(\beta_{g^+}(e, g) + i(\beta_{g^-}(e, g)))} d\nu(g^+) d\nu_1(g^-) da dm$$

where da and dm denote the Haar measures on \mathfrak{a} and M respectively. Hence it induces an AM -invariant measure on $\Gamma \backslash G$ which we also denote by $\hat{m}_{\nu, \nu_1}^{\text{BMS}}$ and call the Bowen-Margulis-Sullivan measure associated to the pair (ν, ν_1) . Note that when ν is of divergence type, ν_1 uniquely exists by Theorem 5.5, and therefore we simply write $\hat{m}_{\nu}^{\text{BMS}} := \hat{m}_{\nu, \nu_1}^{\text{BMS}}$.

Let \mathfrak{D}_{Γ} be the collection of all P° -minimal subsets of $\Gamma \backslash G$ where P° is the identity component of P . For a fixed $\mathcal{E}_0 \in \mathfrak{D}_{\Gamma}$, we set $P_{\Gamma} := \{p \in P : \mathcal{E}_0 p = \mathcal{E}_0\}$. Then P_{Γ} is a finite index co-abelian subgroup of P and is independent of the choice of \mathcal{E}_0 , and moreover the map $P_{\Gamma} \backslash P \rightarrow \mathfrak{D}_{\Gamma}$, $[p] \mapsto \mathcal{E}_0 p$, is bijective [18]. We now present the ergodic decompositions of the Burger-Roblin and Bowen-Margulis-Sullivan measures on $\Gamma \backslash G$:

Theorem 10.4. *Let $\Gamma < G$ be a Zariski dense Π -hypertransverse subgroup. Let ν be a Γ -conformal measure on \mathcal{F} of divergence type. Then*

- (1) $\hat{m}_{\nu}^{\text{BR}} = \sum_{\mathcal{E} \in \mathfrak{D}_{\Gamma}} \hat{m}_{\nu}^{\text{BR}}|_{\mathcal{E}}$ is an N -ergodic decomposition;
- (2) $\hat{m}_{\nu}^{\text{BMS}} = \sum_{\mathcal{E} \in \mathfrak{D}_{\Gamma}} \hat{m}_{\nu}^{\text{BMS}}|_{\mathcal{E}}$ is an A -ergodic decomposition.

In particular, the number of N -ergodic components of $\hat{m}_{\nu}^{\text{BR}}$ and the number of A -ergodic components of $\hat{m}_{\nu}^{\text{BMS}}$ are given by $\#\mathfrak{D}_{\Gamma} = [P : P_{\Gamma}]$.

In [31], Lee-Oh deduced the ergodic decomposition theorem for Π -Anosov subgroups from the ergodicity of NM -action and AM -action on $\Gamma \backslash G$ respectively, which were shown in their another work [32]. The Anosov property was used in order to have

- Π -regularity and Π -antipodality of Γ ;
- the ergodicity of NM -action and the complete conservativity and ergodicity of AM -action on $\Gamma \backslash G$;
- appropriate covering of the limit set to show that the $\mathfrak{a} \times M$ -valued essential subgroup for a Γ -conformal measure ν is the whole $\mathfrak{a} \times M$.

On the other hand, when Γ is Π -hypertransverse, it is Π -regular and Π -antipodal. Moreover, if ν is of divergence type, then we showed that the NM -action on $(\Gamma \backslash G, \hat{m}_{\nu}^{\text{BR}})$ is ergodic in Theorem 1.12 and the complete conservativity and ergodicity of AM -action on $(\Gamma \backslash G, \hat{m}_{\nu}^{\text{BMS}})$ were known ([13], [26], Theorem 5.7). Finally, as we have shown in Section 7, the new

covering of the limit set defined in terms of the visual metric on the Gromov boundary plays an appropriate role to prove that the essential subgroup is full. The deduction for the extended version of the essential subgroup, taking values in $\mathfrak{a} \times M$, can be done in a same way as in [31]. Therefore, Theorem 10.4 can be deduced by the same argument as in [31], with these replacements of the above three items.

REFERENCES

- [1] M. Babillot. On the classification of invariant measures for horosphere foliations on nilpotent covers of negatively curved manifolds. In *Random walks and geometry*, pages 319–335. Walter de Gruyter, Berlin, 2004.
- [2] Y. Benoist. Propriétés asymptotiques des groupes linéaires. *Geom. Funct. Anal.*, 7(1):1–47, 1997.
- [3] Y. Benoist. Propriétés asymptotiques des groupes linéaires. II. In *Analysis on homogeneous spaces and representation theory of Lie groups, Okayama–Kyoto (1997)*, volume 26 of *Adv. Stud. Pure Math.*, pages 33–48. Math. Soc. Japan, Tokyo, 2000.
- [4] G. Besson, G. Courtois, and S. Gallot. Entropies et rigidités des espaces localement symétriques de courbure strictement négative. *Geom. Funct. Anal.*, 5(5):731–799, 1995.
- [5] G. Besson, G. Courtois, and S. Gallot. Minimal entropy and Mostow’s rigidity theorems. *Ergodic Theory Dynam. Systems*, 16(4):623–649, 1996.
- [6] P.-L. Blayac, R. Canary, F. Zhu, and A. Zimmer. Patterson-Sullivan theory for coarse cocycles. *Preprint arXiv:2404.09713*, 2024.
- [7] B. Bowditch. Convergence groups and configuration spaces. In *Geometric group theory down under (Canberra, 1996)*, pages 23–54. de Gruyter, Berlin, 1999.
- [8] H. Bray, R. Canary, L.-Y. Kao, and G. Martone. Counting, equidistribution and entropy gaps at infinity with applications to cusped Hitchin representations. *J. Reine Angew. Math.*, 791:1–51, 2022.
- [9] M. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [10] M. Burger. Horocycle flow on geometrically finite surfaces. *Duke Math. J.*, 61(3):779–803, 1990.
- [11] R. Canary, T. Zhang, and A. Zimmer. Cusped Hitchin representations and Anosov representations of geometrically finite Fuchsian groups. *Adv. Math.*, 404:Paper No. 108439, 67, 2022.
- [12] R. Canary, T. Zhang, and A. Zimmer. Patterson-Sullivan measures for relatively Anosov groups. *Preprint arXiv:2308.04023*, 2023.
- [13] R. Canary, T. Zhang, and A. Zimmer. Patterson-Sullivan measures for transverse subgroups. *Preprint, arXiv:2304.11515*, 2023. To appear in *J. Mod. Dyn.*
- [14] M. Coornaert and A. Papadopoulos. *Symbolic dynamics and hyperbolic groups*, volume 1539 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1993.
- [15] F. Dal’Bo and I. Kim. A criterion of conjugacy for Zariski dense subgroups. *C. R. Acad. Sci. Paris Sér. I Math.*, 330(8):647–650, 2000.
- [16] F. Guéritaud, O. Guichard, F. Kassel, and A. Wienhard. Anosov representations and proper actions. *Geom. Topol.*, 21(1):485–584, 2017.
- [17] O. Guichard and A. Wienhard. Anosov representations: domains of discontinuity and applications. *Invent. Math.*, 190(2):357–438, 2012.
- [18] Y. Guivarc’h and A. Raugi. Actions of large semigroups and random walks on isometric extensions of boundaries. *Ann. Sci. École Norm. Sup. (4)*, 40(2):209–249, 2007.

- [19] G. A. Hedlund. Fuchsian groups and mixtures. *Ann. of Math. (2)*, 40(2):370–383, 1939.
- [20] I. Kapovich and N. Benakli. Boundaries of hyperbolic groups. In *Combinatorial and geometric group theory (New York, 2000/Hoboken, NJ, 2001)*, volume 296 of *Contemp. Math.*, pages 39–93. Amer. Math. Soc., Providence, RI, 2002.
- [21] M. Kapovich, B. Leeb, and J. Porti. Anosov subgroups: dynamical and geometric characterizations. *Eur. J. Math.*, 3(4):808–898, 2017.
- [22] D. M. Kim and H. Oh. Conformal measure rigidity for representations via self-joinings. *arXiv preprint arXiv:2302.03539*, 2023.
- [23] D. M. Kim and H. Oh. Rigidity of Kleinian groups via self-joinings. *Invent. Math.*, 234(3):937–948, 2023.
- [24] D. M. Kim and H. Oh. Rigidity of Kleinian groups via self-joinings: measure theoretic criterion. *arXiv preprint arXiv:2302.03552*, 2023.
- [25] D. M. Kim, H. Oh, and Y. Wang. Ergodic dichotomy for subspace flows in higher rank. *arXiv preprint arXiv:2310.19976*, 2023.
- [26] D. M. Kim, H. Oh, and Y. Wang. Properly discontinuous actions, Growth indicators and Conformal measures for transverse subgroups. *arXiv preprint arXiv:2306.06846*, 2023.
- [27] I. Kim. Length spectrum in rank one symmetric space is not arithmetic. *Proc. Amer. Math. Soc.*, 134(12):3691–3696, 2006.
- [28] F. Labourie. Anosov flows, surface groups and curves in projective space. *Invent. Math.*, 165(1):51–114, 2006.
- [29] O. Landesberg, M. Lee, E. Lindenstrauss, and H. Oh. Horospherical invariant measures and a rank dichotomy for Anosov groups. *J. Mod. Dyn.*, 19:331–362, 2023.
- [30] O. Landesberg and E. Lindenstrauss. On Radon measures invariant under horospherical flows on geometrically infinite quotients. *Int. Math. Res. Not. IMRN*, (15):11602–11641, 2022.
- [31] M. Lee and H. Oh. Ergodic decompositions of geometric measures on Anosov homogeneous spaces. *arXiv:2010.11337*, 2020. To appear in *Isr. J. Math.*
- [32] M. Lee and H. Oh. Invariant measures for horospherical actions and Anosov groups. *Int. Math. Res. Not. IMRN*, (19):16226–16295, 2023.
- [33] G. Mostow. *Strong rigidity of locally symmetric spaces*. Annals of Mathematics Studies, No. 78. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1973.
- [34] G. Prasad. Strong rigidity of \mathbf{Q} -rank 1 lattices. *Invent. Math.*, 21:255–286, 1973.
- [35] J.-F. Quint. Mesures de Patterson-Sullivan en rang supérieur. *Geom. Funct. Anal.*, 12(4):776–809, 2002.
- [36] T. Roblin. Ergodicité et équidistribution en courbure négative. *Mém. Soc. Math. Fr. (N.S.)*, (95):vi+96, 2003.
- [37] A. Sambarino. A report on an ergodic dichotomy. *Ergodic Theory Dynam. Systems*, 44(1):236–289, 2024.
- [38] K. Schmidt. *Cocycles on ergodic transformation groups*. Macmillan Lectures in Mathematics, Vol. 1. Macmillan Co. of India, Ltd., Delhi, 1977.
- [39] D. Sullivan. The density at infinity of a discrete group of hyperbolic motions. *Inst. Hautes Études Sci. Publ. Math.*, (50):171–202, 1979.
- [40] D. Sullivan. Discrete conformal groups and measurable dynamics. *Bull. Amer. Math. Soc. (N.S.)*, 6(1):57–73, 1982.
- [41] P. Tukia. A rigidity theorem for Möbius groups. *Invent. Math.*, 97(2):405–431, 1989.
- [42] P. Tukia. The Poincaré series and the conformal measure of conical and Myrberg limit points. *J. Anal. Math.*, 62:241–259, 1994.
- [43] C. Yue. Mostow rigidity of rank 1 discrete groups with ergodic Bowen-Margulis measure. *Invent. Math.*, 125(1):75–102, 1996.

- [44] F. Zhu and A. Zimmer. Relatively Anosov representations via flows I: theory. *Preprint arXiv:2207.14737*, 2022.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06511
Email address: `dongryul.kim@yale.edu`