

Lecture 13. Refs: II. 5 in Hilton, Stammbach

1) Objects representing functors.

2) Products & coproducts.

"A course in Homological Algebra"

CTM 4.

1.0) Recap: Considered Hom functors: $X \in \text{Ob}(\mathcal{C}) \rightsquigarrow F_X: \mathcal{C} \rightarrow \text{Sets}$

$F_X(Y) = \text{Hom}_{\mathcal{C}}(X, Y)$. We've produced a map

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(X', X) & \xrightarrow{\quad} & \text{Hom}_{\text{Fun}}(F_X, F_{X'}) \\ \downarrow \psi & \downarrow \psi & \\ g & \longmapsto & \psi g \\ \downarrow \psi & \downarrow \psi & \\ \psi g & \xrightarrow{\quad} & \psi g \\ \downarrow \psi & \downarrow \psi & \\ \psi g & \xrightarrow{\quad} & \psi g \end{array}$$

Yoneda lemma: the map $g \mapsto \psi g$ is a bijection.

Also: $\psi g \circ \psi g' = \psi g' g$.

Application: If $F = F_X, G = F_{X'}$, then can compute $\text{Hom}_{\text{Fun}}(F, G)$ (a set), $\text{End}_{\text{Fun}}(F)$, as a monoid:

$\cong \text{End}_{\mathcal{C}}(X)$ as a set, compositions are in opposite order (see example in the end of Lec 12)

Rem*: If \mathcal{C} is a small category $\rightsquigarrow \text{Fun}(\mathcal{C}, \text{Sets})$, then $X \mapsto F_X$ realizes \mathcal{C}^{app} as a full subcategory in $\text{Fun}(\mathcal{C}, \text{Sets})$

from Yoneda

1.1) Objects representing functors.

Definition: Let $F: \mathcal{C} \rightarrow \text{Sets}$. We say $X \in \text{Ob}(\mathcal{C})$ represents F if F is isomorphic to F_X .

Example (from Lec 12): Forgetful functor $\text{Groups} \rightarrow \text{Sets}$

is represented by \mathbb{Z} .

A representing object may fail to exist.

Lemma: An object, X , representing $F: \mathcal{C} \rightarrow \text{Sets}$ is unique (up to isomorphism) if it exists.

Proof: Two representing objects $X, X': F_X \xrightarrow{\sim} F \xleftarrow{\sim} F_{X'}$,
so $\eta: F_X \xrightarrow{\sim} F_{X'} \xleftarrow{\text{Yoneda}} X' \xrightarrow{g} X$ (w. $\eta = \eta \circ g$).

Since $g \mapsto \eta \circ g$ is compatible w. comp'n's, η is invertible \Rightarrow
 g is invertible $\Rightarrow X, X'$ are isomorphic. \square

2) Products & coproducts.

2.1) Definition of product.

General constr'n: $F_1, F_2: \mathcal{C} \rightarrow \text{Sets}$. Can take direct products of sets $\leadsto F_1 \times F_2: \mathcal{C} \rightarrow \text{Sets}$

for $Y \in \text{Ob}(\mathcal{C}) \leadsto F_1 \times F_2(Y) = F_1(Y) \times F_2(Y)$

for $Y \xrightarrow{f} Y' \leadsto F_1 \times F_2(f) = F_1(f) \times F_2(f): F_1(Y) \times F_2(Y) \rightarrow F_1(Y') \times F_2(Y')$


Definition: Let $X_1, X_2 \in \text{Ob}(\mathcal{C})$. By their product we mean an object, denoted $X_1 \times X_2$, representing the functor $F_{X_1}^{\text{opp}} \times F_{X_2}^{\text{opp}}$ (i.e. $Y \mapsto \text{Hom}_{\mathcal{C}}(Y, X_1) \times \text{Hom}_{\mathcal{C}}(Y, X_2)$ on objects).

Note: $X_1 \times X_2$ may fail to exist.

Exercise (on Lemma): $X_1 \times X_2$ is isom'ic to $X_2 \times X_1$.

Alternative definition (via universal property): Let $X_1, X_2 \in \text{Ob}(\mathcal{C})$

By the product of X_1 & X_2 we mean (X, π_1, π_2) w. $X \in \mathcal{Ob}(\mathcal{C})$,
 $\pi_i \in \text{Hom}_{\mathcal{C}}(X, X_i)$ ($i=1,2$) s.t. $\forall Y \in \mathcal{Ob}(\mathcal{C}), \psi_i \in \text{Hom}_{\mathcal{C}}(Y, X_i)$
 $\exists! \psi \in \text{Hom}_{\mathcal{C}}(Y, X)$ s.t. the following is commutative:



Lemma: The two definitions are equivalent.

Proof: An object constructed in Altern. def'n satisfies Definition:

Let (X, π_1, π_2) be as above. We need a functor isomorphism

$$\eta: \text{Hom}_{\mathcal{C}}(\cdot, X) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(\cdot, X_1) \times \text{Hom}_{\mathcal{C}}(\cdot, X_2)$$

• Construct $\eta_Y: \text{Hom}_{\mathcal{C}}(Y, X) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(Y, X_1) \times \text{Hom}_{\mathcal{C}}(Y, X_2)$
 $\psi \longmapsto (\pi_1 \circ \psi, \pi_2 \circ \psi)$

η_Y is a bijection b/c of the universal property.

• Check η is a functor morphism, i.e. $\forall Y' \xrightarrow{f} Y$

the following diagram commutes:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(Y, X) & \xrightarrow{\eta_Y: \psi \mapsto (\pi_1 \circ \psi, \pi_2 \circ \psi)} & \text{Hom}_{\mathcal{C}}(Y, X_1) \times \text{Hom}_{\mathcal{C}}(Y, X_2) \\ \downarrow \psi \mapsto \psi \circ f & & \downarrow (\psi_1, \psi_2) \mapsto (\psi_1 \circ f, \psi_2 \circ f) \\ \text{Hom}_{\mathcal{C}}(Y', X) & \xrightarrow{\eta_{Y'}: \psi' \mapsto (\pi_1 \circ \psi', \pi_2 \circ \psi')} & \text{Hom}_{\mathcal{C}}(Y', X_1) \times \text{Hom}_{\mathcal{C}}(Y', X_2) \end{array}$$

- exercise.

Now: Definition \rightsquigarrow Alternative definition:

Note $F_1 \times F_2$ admits a functor morphism $\pi^i: F_1 \times F_2 \rightarrow F_i$

π^i is the projection $F_1(Y) \times F_2(Y) \rightarrow F_i(Y)$ (exercise). Now we have

$$\eta: \text{Hom}_E(\cdot, X_1 \times X_2) \longrightarrow \text{Hom}_E(\cdot, X_1) \times \text{Hom}_E(\cdot, X_2) \rightsquigarrow$$

$$\pi^i \circ \eta: \text{Hom}_E(\cdot, X_1 \times X_2) \longrightarrow \text{Hom}_E(\cdot, X_i). \text{ By Yoneda,}$$

$$\pi^i \circ \eta = \eta^{\pi^i} \text{ for the unique } \pi_i \in \text{Hom}_E(X_1 \times X_2, X_i) \text{ (} i=1,2 \text{):}$$

$(\pi^i \circ \eta)_Y(\psi) = \pi_i \circ \psi$. ($\forall \psi \in \text{Hom}_E(Y, X_1 \times X_2)$). Since η is a functor isomorphism $\Rightarrow \eta_Y = ((\pi^1 \circ \eta)_Y, (\pi^2 \circ \eta)_Y)$ is a bijection

$$\text{Hom}_E(Y, X_1 \times X_2) \xrightarrow{\sim} \text{Hom}_E(Y, X_1) \times \text{Hom}_E(Y, X_2).$$


$$\underbrace{\psi}_{\eta} \longmapsto (\underbrace{\pi_1 \circ \psi}_{\pi^1 \circ \eta}, \underbrace{\pi_2 \circ \psi}_{\pi^2 \circ \eta}) (= \eta_Y(\psi))$$

So we recover the universal property. \square

2.2) Coproduct := product in \mathcal{C}^{opp} . Notation: $X_1 * X_2$

Universal property: have $X_i \xrightarrow{\iota_i} X_1 * X_2$ s.t. $\forall Y \in \text{Obj}(\mathcal{C})$, $X_i \xrightarrow{\varphi_i} Y, i=1,2 \quad \exists! X_1 * X_2 \xrightarrow{\varphi} Y$ s.t. the following is

comm'ive:



Eqv.: $X_1 * X_2$ represents $\text{Hom}_E(X_1, \cdot) \times \text{Hom}_E(X_2, \cdot)$.

2.3) Examples: Products in Sets, Groups, Rings, A-Mod etc:

product = direct product.

Coproducts:

In Sets, coproduct = disjoint union.

In A -Mod, coproduct = direct sum (see Prob 7 in HW1)

In general alg'c cat's can talk about objects defined by generators & relations:

Let X_1 be given by generators $a_i, (i \in I)$, w. relations $F_\alpha, \alpha \in A$; let X_2 have generators $b_j, (j \in J)$ w. relations $G_\beta, \beta \in B$

Then $X_1 * X_2$ is generated by the union: $a_i (i \in I), b_j (j \in J)$ w. relations F_α (on a_i 's) & G_β (on b_j 's) - also the union.

2.4) Remarks.

- can talk about (co)products indexed by an arbitrary set, I (rather than $\{1,2\}$), for example, direct sums & products in A -Mod.

- products may fail to exist: Let \mathcal{C} be the full subcat. in \mathbb{F} -Vect w. objects being odd-dimensional vector spaces. For any two objects in \mathcal{C} , there's no product (in \mathcal{C}).

2.5) Initial/terminal objects in cat's.

Definition: Let \mathcal{C} be a category. Say $X \in \text{Ob}(\mathcal{C})$ is:

• initial if $\forall Y \in \text{Ob}(\mathcal{C}) \Rightarrow \text{Hom}_{\mathcal{C}}(X, Y)$ consists of 1 element.

• terminal $\dots \dots \dots \text{Hom}_{\mathcal{C}}(Y, X) \dots \dots \dots$

Initial in $\mathcal{C} = \text{terminal in } \mathcal{C}^{\text{opp}}$

Examples: • In A -Mod, $\{0\}$ is both initial & terminal

• In Rings: initial object: \mathbb{Z}

terminal object: $\{0\}$.

Exercise: For $X_1, X_2 \in \text{Ob}(\mathcal{C}) \rightsquigarrow$ new cat. $\mathcal{L}(X_1, X_2)$

Ob: triples $\{(Y, \psi_1, \psi_2) \mid Y \in \text{Ob}(\mathcal{C}), \psi_i \in \text{Hom}_{\mathcal{C}}(Y, X_i), i=1,2\}$

$\text{Hom}((Y, \psi_1, \psi_2), (Y', \psi'_1, \psi'_2)) := \{\varphi \in \text{Hom}_{\mathcal{C}}(Y, Y') \mid \psi_i = \psi'_i \circ \varphi, i=1,2\}$

Composition of morphisms is restricted from \mathcal{C} .

Persuade yourself that $\mathcal{L}(X_1, X_2)$ is a category &
 $(X_1 \times X_2, \pi_1, \pi_2)$ is a terminal object (if exists)