

## Lecture 4: rings, ideals & modules, IV.

- 1) Constructions w. modules.
- 2) Submodules & quotient modules.
- 3) Finitely generated & free modules.

References: [AM], Chapter 2, Sections 2,3,5.

### 1) Construction with modules & homomorphisms: Hom module

Let  $A$  be a commutative ring &  $M, N$  be  $A$ -modules. Define the set  $\text{Hom}_A(M, N) := \{A\text{-linear maps } M \rightarrow N\}$

Claim:  $\text{Hom}_A(M, N)$  is  $A$ -module w.r.t. the point-wise operations:  
for  $\psi, \psi' \in \text{Hom}_A(M, N)$ ,  $a \in A$

$$[\psi + \psi'](m) := \psi(m) + \psi'(m) \in N \quad (\leadsto \psi + \psi' \text{ is a map } M \rightarrow N)$$

$$[a\psi](m) := a\psi(m) \in N$$

Lemma: 1)  $\psi + \psi'$ ,  $a\psi$  are  $A$ -linear maps.

2) The operations  $+$ ,  $\cdot$  turn  $\text{Hom}_A(M, N)$  into  $A$ -module.

Partial proof:  $[a\psi](bm) = b[a\psi](m)$  - part of linearity for  $a\psi$ .

$$[a\psi](bm) = a(\psi(bm)) = ab\psi(m) = [ab = ba] = b(a\psi(m)) = b[a\psi](m).$$

Rest of proof is an exercise.  $\square$

Example: 1) Let  $M = A$ . Then  $\text{Hom}_A(A, N) \xrightarrow{\sim} N$  as  $A$ -modules.

Namely, we have a map  $\text{Hom}_A(A, N) \rightarrow N$ ,  $\psi \mapsto \psi(1)$ . It's  $A$ -linear & has inverse: for  $n \in N$  define the map  $\psi_n: A \rightarrow N$ ,  $a \mapsto an$ . The inverse is  $n \mapsto \psi_n$ . To check details (that these maps are well-defined & mutually inverse) is an **exercise**.  $\square$

$$2) \text{Hom}_A(M_1 \oplus M_2, N) \xrightarrow{\sim} \text{Hom}_A(M_1, N) \times \text{Hom}_A(M_2, N)$$

$$\psi \mapsto (\psi|_{M_1}, \psi|_{M_2}), \text{ where } M_1 \hookrightarrow M_1 \oplus M_2 \text{ via } m_1 \mapsto (m_1, 0), \text{ and similarly for } M_2.$$

Inverse map:  $(\psi_1, \psi_2) \in \text{Hom}_A(M_1, N) \times \text{Hom}_A(M_2, N)$  goes to  $\psi: M_1 \oplus M_2 \rightarrow N$  given by  $\psi(m_1, m_2) := \psi_1(m_1) + \psi_2(m_2)$

**Exercise:** Prove that these two maps are mutually inverse  $A$ -module homomorphisms.

3) There is a direct analog of example 2) for  $M_1 \oplus \dots \oplus M_k$ .

$$\text{E.g. } \text{Hom}_A(A^{\oplus k}, N) \xrightarrow{\sim} \text{Hom}_A(A, N)^{\times k} \xrightarrow{\sim} N^{\times k}$$

$$\psi \longmapsto (\psi(e_1), \dots, \psi(e_k))$$

where  $e_i = (0, \dots, 1, \dots, 0)$  (1 in the  $i$ th place). The inverse map is given by  $\underline{n} = (n_1, \dots, n_k) \mapsto \psi_{\underline{n}}: (a_1, \dots, a_k) \mapsto \sum a_i n_i$ .

**Rem:** Example 2 further generalizes to infinite direct sums:

$$\text{Hom}_A\left(\bigoplus_{i \in I} M_i, N\right) \xrightarrow{\sim} \prod_{i \in I} \text{Hom}_A(M_i, N)$$

The proof is similar to the above and is left as **exercise**.

## 2) Sub & quotient modules.

### 2.1) Submodules:

Let  $A$  be a comm'ive ring.

**Definition:** Let  $M$  be an  $A$ -module; a **submodule** in  $M$  is an abelian subgroup  $N \subset M$  s.t.  $a \in A, n \in N \Rightarrow an \in N$ .

**Rem:**  $N$  has a natural  $A$ -module str'ure.

**Examples:** 0)  $\{0\}, M \subset M$  are submodules.

1)  $A$  is a field (so module = vector space): Submodule = subspace.

2)  $A = \mathbb{Z}$  (so module = abelian group): Submodule = subgroup.

3)  $A = \mathbb{F}[x]$  ( $\mathbb{F}$  is a field). A module  $M = \mathbb{F}$ -vector space w. operator  $X: M \rightarrow M$ . A submodule  $N \subset M$  - subspace s.t.  $X(N) \subseteq N$ . Conversely, every  $X$ -stable subspace is a submodule b/c  $f(x)m = f(X)m$  &  $X(N) \subseteq N \Rightarrow f(X)(N) \subset N$ .

4)  $A$  is any ring,  $M = A$ : submodule = ideal.

### 2.2) Constructions w. submodules.

1)  $\psi: M \rightarrow N$   $A$ -module homom'm:  $\ker \psi \subset M$  &  $\text{im } \psi \subset N$  are submodules, left as **exercise**.

2)  $m_1, \dots, m_k \in M \leadsto \text{Span}_A(m_1, \dots, m_k) := \left\{ \sum_{i=1}^k a_i m_i \mid a_i \in A \right\}$  - this is special case of image:  $\underline{m} = (m_1, \dots, m_k) \leadsto \psi_{\underline{m}}: A^{\oplus k} \rightarrow M$  (see example 3 in Sec 1.2) Then  $\text{Span}_A(m_1, \dots, m_k) = \text{im } \psi_{\underline{m}}$ . Note also that this generalizes the ideal generated by a given collection of elements, Sec 3.1 of Lec 1. More generally, for index set  $I$  &  $m_i \in M$  ( $i \in I$ )  $\leadsto \text{Span}_A(m_i \mid i \in I) = \{\text{finite } A\text{-linear combinations of } m_i\text{'s}\}$ .

3) Sums & intersections:  $M_1, M_2 \subset M$  submodules  
 $M_1 \cap M_2, M_1 + M_2 = \{m_1 + m_2 \mid m_i \in M_i\}$  - submodules.

4) Product w. ideal:  $N \subset M$  submodule,  $I \subset A$  ideal  
 $IN := \left\{ \sum_{i=1}^n a_i n_i \mid a_i \in I, n_i \in N \right\}$  - submodule, **exercise**.  
 (compare to product of ideals in 1.1 of Lecture 2).

**2.3) Quotient modules:**  $M$  is  $A$ -module,  $N \subset M$  submodule  
 $\leadsto$  abelian group  $M/N = \{m+N \mid m \in M\}$  & abelian group homom'm  
 $\pi: M \rightarrow M/N, \pi(m) := m+N$ . Then  $M/N$  has a natural  $A$ -module str'ure. The following is analogous to Proposition in Sec 3.2 of Lecture 1.

**Proposition:** 1) The map  $A \times (M/N) \rightarrow M/N, (a, m+N) \mapsto am+N$  is well-defined ( $am+N$  only depends on  $m+N$  & not on  $m$  itself) and equips  $M/N$  w.  $A$ -module structure.

2) This module structure is unique s.t.  $\pi: M \rightarrow M/N$  is a module homomorphism.

3) (Universal property of  $M/N$  &  $\pi: M \rightarrow M/N$ ) Let  $\psi: M \rightarrow M'$  be  $A$ -module homom'm s.t.  $N \subset \ker \psi$ . Then  $\exists!$  module homom'm  $\underline{\psi}: M/N \rightarrow M'$  s.t. the following diagram is commutative

$$\begin{array}{ccc} M & \xrightarrow{\psi} & M' \\ \pi \downarrow & \searrow & \\ M/N & \xrightarrow{\underline{\psi}} & M' \end{array} \quad \begin{array}{l} \psi \text{ is given by:} \\ \psi(m+N) := \psi(m) \end{array}$$

Proof: exercise.

Remarks:

1) Let  $I \subset A$  be ideal  $\leadsto$  submodule  $IM \subset M \leadsto$  quotient  $M/IM$  is  $A$ -module where  $I$  acts by 0 =  $A/I$ -module; explicitly  $(a+I)(m+IM) = am+IM$  (see Observation II in Sec 2.2 of Lec 3).

2) We have standard "isomorphism theorems":

- for  $\psi: M \rightarrow N$ ,  $A$ -module homom'm, have  $M/\ker \psi \xrightarrow{\sim} \text{im } \psi$  ( $A$ -module isomorphism).

- for submodules  $K \subset N \subset M$ , have  $(M/K)/(N/K) \xrightarrow{\sim} M/N$ .

- for submodules  $N_1, N_2 \subset M$ , have  $N_1/N_1 \cap N_2 \xrightarrow{\sim} (N_1+N_2)/N_2$ .

The reason is that the standard abelian group isomorphisms are also module isomorphisms.

3) There are bijections between:

$$\begin{aligned} & \{ \text{submodules } L \subset M \mid N \subset L \} \\ & \left( L \mapsto \pi(L) = L/N \right) \\ & \left( L \mapsto \pi^{-1}(L) \right) \\ & \{ \text{submodules } \underline{L} \subset M/N \} \end{aligned}$$

We've seen a similar claim for ideals in Sec. 3.2 of Lec. 1.

3) Finitely generated & free modules.

3.1) Finitely generated modules

Definition: Elements  $m_i \in M$  ( $i \in I$ ) are **generators** (a.k.a. **spanning set**) of  $M$  if  $M = \text{Span}_A(m_i \mid i \in I)$ , i.e.  $\forall m \in M$  is  $A$ -linear combination of finite number of  $m_i$ 's.

•  $M$  is **finitely generated** if it has a finite spanning set.

Remarks: 1)  $A^{\oplus I}$  is finitely generated  $\Leftrightarrow I$  is finite.

2) If  $M$  is fin. generated, then so is  $M/N$   $\forall N \subset M$ :  
 $M = \text{Span}_A(m_1, \dots, m_k) \Rightarrow M/N = \text{Span}_A(\pi(m_1), \dots, \pi(m_k)).$

3)  $\underline{m} = (m_1, \dots, m_k) \rightsquigarrow \psi_{\underline{m}}: A^{\oplus k} \rightarrow M, \psi_{\underline{m}}(a_1, \dots, a_k) = \sum_{i=1}^k a_i m_i$ .  
 $M = \text{Span}_A(m_1, \dots, m_k) \Leftrightarrow \psi_{\underline{m}}$  is surj'ive  $\Rightarrow M \cong A^{\oplus k} / \ker \psi_{\underline{m}}$ .  
 So: fin. gen'd modules = quotients of  $A^{\oplus k}$  for some  $k \in \mathbb{N}_0$ .

### 3.2) Free modules

Let  $M$  be an  $A$ -module.

**Definition:** Elements  $m_i, i \in I$ , form a **basis** in  $M$  if  $\forall m \in M$  is uniquely written as  $A$ -linear combination of  $m_i, i \in I$ .  
•  $M$  is **free** if it has a basis.

**Examples:** 1) For any set  $I$ ,  $A^{\oplus I}$  is free, for a basis can take coordinate vectors  $e_i = (0, \dots, \underset{\substack{\uparrow \\ \text{ith place}}}{1}, \dots, 0, \dots)$ ,  $i \in I$ .

2) If  $A$  is field, then every module (a.k.a. vector space) is free. If  $A$  is not a field, there are non-free modules: let  $J \subset A$  be ideal,  $J \neq \{0\}$ ,  $A \Rightarrow A/J$  is not free (over  $A$ ).

Indeed, for any vector  $e$  in a basis we must have  $ae \neq 0 \forall a \in A$ . But for any  $e \in A/J$  we have  $ae = 0 \forall a \in J$ .

**Remarks** 1) Every free module is isomorphic to  $A^{\oplus I}$  for some set  $I$ : choose basis  $m_i \in M$  ( $i \in I$ ):  $\varphi_m: A^{\oplus I} \xrightarrow{\sim} M$ . Moreover, Problem 5 in HW1 shows that  $I$  is uniquely determined by  $M$  (up to bijection of sets)

2) As for vector spaces, any homomorphism  $\varphi: A^{\oplus k} \rightarrow A^{\oplus \ell}$  is given by multiplication by an  $\ell \times k$ -matrix (where elements of  $A^{\oplus k}, A^{\oplus \ell}$  are viewed as column vectors) & the matrix is uniquely determined by  $\varphi$ .

7