

## INVARIANT THEORY, HW3

**Problem 1, 7pts.** This problem describes the algebra of invariants  $\mathbb{C}[V]^T$  under a faithful action of a torus  $T$  on a vector space  $V$ . Recall that  $V$  has an eigen-basis  $v_1, \dots, v_n$ , let  $\chi_1, \dots, \chi_n$  be the corresponding eigen-characters of  $T$ . Let  $\tilde{T} \subset \text{GL}(V)$  be the maximal torus of all operators diagonal in the basis  $v_1, \dots, v_n$ . So  $T \subset \tilde{T}$ . Note that  $\tilde{T}/T$  is a torus naturally acting on  $\mathbb{C}[V]^T$  and the character lattice  $\mathfrak{X}(\tilde{T}/T)$  naturally embeds into  $\mathfrak{X}(\tilde{T}) = \mathbb{Z}^n$ . Let  $\mathcal{M}$  denote the submonoid  $\{(m_1, \dots, m_n) \in \mathbb{Z}_{\geq 0}^n \mid \sum_{i=1}^n m_i \chi_i = 0\} \subset \mathbb{Z}_{\geq 0}^n$ .

a, 3pts) Show that each eigenspace of  $\tilde{T}/T$  in  $\mathbb{C}[V]^T$  has dimension 0 or 1 and the dimension is equal to 1 iff the corresponding eigen-character lies in  $\mathcal{M}$ . For  $\psi \in \mathcal{M}$  let  $f_\psi$  denote an eigen-vector for  $\psi$  in  $\mathbb{C}[V]^T$ .

b, 2pts) Show that elements  $\psi_1, \dots, \psi_k$  generate the monoid  $\mathcal{M}$  iff the polynomials  $f_{\psi_1}, \dots, f_{\psi_k}$  generate the algebra  $\mathbb{C}[V]^T$ . Deduce that  $\mathcal{M}$  is a finitely generated monoid.

c, 2pts) Show that the relations of the form  $f_{\psi_1}^{s_1} \dots f_{\psi_k}^{s_k} - f_{\psi_1}^{r_1} \dots f_{\psi_k}^{r_k} = 0$  with  $s_i, r_i \geq 0$  and  $\sum_{i=1}^k (s_i - r_i) \psi_i = 0$  generate the ideal of relations between the generators  $f_{\psi_1}, \dots, f_{\psi_k}$ .<sup>1</sup>

**Problem 2, 7pts.** This problem studies the subvarieties of nilpotent elements (=elements whose orbit closures contain 0) for natural actions of classical groups. Use the Hilbert-Mumford theorem<sup>2</sup> to prove the following claims.

a, 2pts) Let  $G = \text{GL}(U)$  act on  $V := U^{\oplus k} \oplus U^{*\oplus \ell}$  in a natural way. Show that  $(u_1, \dots, u_k, u^1, \dots, u^\ell)$  is nilpotent iff  $\langle u_i, u^j \rangle = 0$  for all  $i, j$ .

b) 2pts) Let  $G = \text{SL}(U)$  acts on  $U^{\oplus k}$ . Then  $(u_1, \dots, u_k)$  is nilpotent iff  $u_1, \dots, u_k$  do not span  $U$ .

c, 3pts) Let  $U$  be a finite dimensional space equipped with an orthogonal form  $(\cdot, \cdot)$ . Consider the natural action of the orthogonal group  $G = \text{O}(U)$  on  $V = U^{\oplus k}$ . Then  $(u_1, \dots, u_k)$  is nilpotent iff  $(u_i, u_j) = 0$  for all  $i, j$ . State and prove the corresponding claim for the symplectic group.

<sup>1</sup>Of course, there is a more economical set of relations.

<sup>2</sup>Solutions based on computing the algebras of invariants do not count