

# Long-term dynamics of nonlinear wave equations

W. Schlag (University of Chicago)

ICM, Seoul 2014

## Wave maps

Let  $(M, g)$  be a Riemannian manifold, and  $u : \mathbb{R}_{t,x}^{1+d} \rightarrow M$  smooth.

Wave maps defined by Lagrangian

$$\mathcal{L}(u, \partial_t u) = \int_{\mathbb{R}_{t,x}^{1+d}} \frac{1}{2} (-|\partial_t u|_g^2 + \sum_{j=1}^d |\partial_j u|_g^2) dt dx$$

Critical points  $\mathcal{L}'(u, \partial_t u) = 0$  satisfy “manifold-valued wave equation”.  
 $M \subset \mathbb{R}^N$  embedded, this equation is

$$\square u \perp T_u M \text{ or } \square u = A(u)(\partial u, \partial u),$$

$A$  being the second fundamental form.

For example,  $M = \mathbb{S}^{n-1}$ , then

$$\square u = u(|\partial_t u|^2 - |\nabla u|^2)$$

Note: Nonlinear wave equation, null-form! Harmonic maps are solutions.

## Wave maps

Intrinsic formulation:  $D^\alpha \partial_\alpha u = \eta^{\alpha\beta} D_\beta \partial_\alpha u = 0$ , in coordinates

$$-\partial_{tt} u^i + \Delta u^i + \Gamma_{jk}^i(u) \partial_\alpha u^j \partial^\alpha u^k = 0$$

$\eta = (-1, 1, 1, \dots, 1)$  Minkowski metric

- Similarity with geodesic equation:  $u = \gamma \circ \varphi$  is a wave map provided  $\square \varphi = 0$ ,  $\gamma$  a geodesic.

- Energy conservation:  $E(u, \partial_t u) = \int_{\mathbb{R}^d} (|\partial_t u|_g^2 + \sum_{j=1}^d |\partial_j u|_g^2) dx$  is conserved in time.

- Cauchy problem:

$$\square u = A(u)(\partial^\alpha u, \partial_\alpha u), \quad (u(0), \partial_t u(0)) = (u_0, u_1)$$

smooth data. Does there exist a smooth local or global-in-time solution?

**Local:** Yes. **Global:** depends on the dimension of Minkowski space and the geometry of the target.

## Criticality and dimension

If  $u(t, x)$  is a wave map, then so is  $u(\lambda t, \lambda x)$ ,  $\forall \lambda > 0$ .

Data in the Sobolev space  $\dot{H}^s \times \dot{H}^{s-1}(\mathbb{R}^d)$ . For which  $s$  is this space invariant under the natural scaling? Answer:  $s = \frac{d}{2}$ .

Scaling of the energy:  $u(t, x) \mapsto \lambda^{\frac{d-2}{2}} u(\lambda t, \lambda x)$  same as  $\dot{H}^1 \times L^2$ .

- **Subcritical case:**  $d = 1$  the natural scaling is associated with less regularity than that of the conserved energy. Expect global existence. Logic: local time of existence only depends on energy of data, which is preserved.
- **Critical case:**  $d = 2$ . Energy keeps the balance with the natural scaling of the equation. For  $\mathbb{S}^2$  can have finite-time blowup, whereas for  $\mathbb{H}^2$  have global existence. Krieger-S.-Tataru 06, Krieger-S. 09, Rodnianski-Raphael 09, Sterbenz-Tataru 09.
- **Supercritical case:**  $d \geq 3$ . Poorly understood. Self-similar blowup  $Q(r/t)$  for sphere as target, Shatah 80s. Also negatively curved manifolds possible in high dimensions: Cazenave, Shatah, Tahvildar-Zadeh 98.

## A nonlinear defocusing Klein-Gordon equation

Consider in  $\mathbb{R}_{t,x}^{1+3}$

$$\square u + u + u^3 = 0, \quad (u(0), \dot{u}(0)) = (f, g) \in \mathcal{H} := H^1 \times L^2(\mathbb{R}^3)$$

Conserved energy

$$E(u, \dot{u}) = \int_{\mathbb{R}^3} \left( \frac{1}{2} |\dot{u}|^2 + \frac{1}{2} |\nabla u|^2 + \frac{1}{2} |u|^2 + \frac{1}{4} |u|^4 \right) dx$$

With  $S(t)$  the linear propagator of  $\square + 1$  we have

$$\vec{u}(t) = (u, \dot{u})(t) = S(t)(f, g) - \int_0^t S(t-s)(0, u^3(s)) ds$$

whence by a simple energy estimate,  $I = (0, T)$

$$\begin{aligned} \|\vec{u}\|_{L^\infty(I; \mathcal{H})} &\lesssim \|(f, g)\|_{\mathcal{H}} + \|u^3\|_{L^1(I; L^2)} \lesssim \|(f, g)\|_{\mathcal{H}} + \|u\|_{L^3(I; L^6)}^3 \\ &\lesssim \|(f, g)\|_{\mathcal{H}} + T \|\vec{u}\|_{L^\infty(I; \mathcal{H})}^3 \end{aligned}$$

Contraction for small  $T$  implies local wellposedness for  $\mathcal{H}$  data.

## Defocusing NLKG3

$T$  depends only on  $\mathcal{H}$ -size of data. From energy conservation we obtain **global existence** by time-stepping.

**Scattering (as in linear theory):**  $\|\vec{u}(t) - \vec{v}(t)\|_{\mathcal{H}} \rightarrow 0$  as  $t \rightarrow \infty$  where  $\square v + v = 0$  energy solution.

$$\vec{v}(0) := \vec{u}(0) - \int_0^\infty S(-s)(0, u^3)(s) ds \quad \text{provided } \|u^3\|_{L_t^1 L_x^2} < \infty$$

**Strichartz estimate** uniformly in intervals  $I$

$$\|\vec{u}\|_{L^\infty(I; \mathcal{H})} + \|u\|_{L^3(I; L^6)} \lesssim \|(f, g)\|_{\mathcal{H}} + \|u\|_{L^3(I; L^6)}^3$$

**Small data scattering:**  $\|\vec{u}\|_{L^3(I; L^6)} \lesssim \|(f, g)\|_{\mathcal{H}} \ll 1$  for all  $I$ . So  $I = \mathbb{R}$  as desired.

**Large data scattering valid; induction on energy, concentration compactness** (Bourgain, Bahouri-Gerard, Kenig-Merle).

## Scattering blueprint

Let  $\vec{u}$  be nonlinear solution with data  $(u_0, u_1) \in \mathcal{H}$ . **Forward scattering set**

$$\mathcal{S}_+ = \{(u_0, u_1) \in \mathcal{H} \mid \vec{u}(t) \text{ exists globally, scatters as } t \rightarrow +\infty\}$$

We claim that  $\mathcal{S}_+ = \mathcal{H}$ . This is proved via the following outline:

- (Small data result):  $\|(u_0, u_1)\|_{\mathcal{H}} < \varepsilon$  implies  $(u_0, u_1) \in \mathcal{S}_+$
- (Concentration Compactness): **If scattering fails**, i.e., if  $\mathcal{S}_+ \neq \mathcal{H}$ , then construct  $\vec{u}_*$  of **minimal energy**  $E_* > 0$  for which  $\|u_*\|_{L_t^3 L_x^6} = \infty$ . There exists  $x(t)$  so that the trajectory

$$K_+ = \{\vec{u}_*(\cdot - x(t), t) \mid t \geq 0\}$$

is **pre-compact** in  $\mathcal{H}$ .

- (Rigidity Argument): If a forward global evolution  $\vec{u}$  has the property that  $K_+$  pre-compact in  $\mathcal{H}$ , then  $u \equiv 0$ .

Kenig-Merle 2006, Bahouri-Gérard decomposition 1998; Merle-Vega.

## Bahouri-Gérard: symmetries vs. dispersion

Let  $\{u_n\}_{n=1}^\infty$  free Klein-Gordon solutions in  $\mathbb{R}^3$  s.t.

$$\sup_n \|\vec{u}_n\|_{L_t^\infty \mathcal{H}} < \infty$$

$\exists$  free solutions  $v^j$  bounded in  $\mathcal{H}$ , and  $(t_n^j, x_n^j) \in \mathbb{R} \times \mathbb{R}^3$  s.t.

$$u_n(t, x) = \sum_{1 \leq j < J} v^j(t + t_n^j, x + x_n^j) + w_n^J(t, x)$$

satisfies  $\forall j < J$ ,  $\vec{w}_n^J(-t_n^j, -x_n^j) \rightarrow 0$  in  $\mathcal{H}$  as  $n \rightarrow \infty$ , and


- $\lim_{n \rightarrow \infty} (|t_n^j - t_n^k| + |x_n^j - x_n^k|) = \infty \forall j \neq k$
- dispersive errors  $w_n^k$  vanish asymptotically:

$$\lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|w_n^J\|_{(L_t^\infty L_x^p \cap L_t^3 L_x^6)(\mathbb{R} \times \mathbb{R}^3)} = 0 \quad \forall 2 < p < 6$$

- orthogonality of the energy:

$$\|\vec{u}_n\|_{\mathcal{H}}^2 = \sum_{1 \leq j < J} \|\vec{v}^j\|_{\mathcal{H}}^2 + \|\vec{w}_n^J\|_{\mathcal{H}}^2 + o(1)$$

## Profiles and Strichartz sea



We can extract further profiles from the Strichartz sea if  $w_n^4$  does not vanish as  $n \rightarrow \infty$  in a suitable sense. In the **radial case** this means  $\lim_{n \rightarrow \infty} \|w_n^4\|_{L_t^\infty L_x^p(\mathbb{R}^3)} > 0$ .

## Lorentz transformations

$$\begin{bmatrix} t' \\ x'_1 \\ x'_2 \\ x'_3 \end{bmatrix} = \begin{bmatrix} \cosh \alpha & \sinh \alpha & 0 & 0 \\ \sinh \alpha & \cosh \alpha & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} t \\ x_1 \\ x_2 \\ x_3 \end{bmatrix}$$




Figure : Causal structure of space-time

## Further remarks on Bahouri-Gérard

- Noncompact symmetry groups: space-time translations and Lorentz transforms.

Compact symmetry groups: Rotations

Lorentz transforms do not appear in the profiles: Energy bound compactifies them.

- Dispersive error  $w_n^j$  is not an energy error!
- In the radial case only need time translations

## The focusing NLKG equation

The **focusing** NLKG

$$\square u + u = \partial_{tt} u - \Delta u + u = u^3$$

has **indefinite conserved energy**


$$E(u, \dot{u}) = \int_{\mathbb{R}^3} \left( \frac{1}{2} |\dot{u}|^2 + \frac{1}{2} |\nabla u|^2 + \frac{1}{2} |u|^2 - \frac{1}{4} |u|^4 \right) dx$$

- Local wellposedness for  $H^1 \times L^2(\mathbb{R}^3)$  data
- Small data: **global existence and scattering**
- **Finite time blowup**  $u(t) = \sqrt{2}(T-t)^{-1}(1 + o(1))$  as  $t \rightarrow T-$   
Cutoff to a cone using finite propagation speed to obtain finite energy solution.
- **stationary solutions**  $-\Delta \varphi + \varphi = \varphi^3$ , ground state  $Q(r) > 0$

## Payne-Sattinger theory; saddle structure of energy near $Q$

Criterion: finite-time blowup/global existence?

Yes, provided the energy is less than the ground state energy Payne-Sattinger 1975.



$$J(\varphi) = \int_{\mathbb{R}^3} \left( \frac{1}{2} |\nabla \varphi|^2 + \frac{1}{2} |\varphi|^2 - \frac{1}{4} |\varphi|^4 \right) dx$$

$$K(\varphi) = \int_{\mathbb{R}^3} (|\nabla \varphi|^2 + |\varphi|^2 - |\varphi|^4) dx$$

**Uniqueness of  $Q$  is the foundation!**

## Payne-Sattinger theory

$j_\varphi(\lambda) := J(e^\lambda \varphi)$ ,  $\varphi \neq 0$  fixed.




Figure : Payne-Sattinger well

Normalize so that  $\lambda_* = 0$ . Then  $\partial_\lambda j_\varphi(\lambda)|_{\lambda=\lambda_*} = K(\varphi) = 0$ .

“Trap” the solution in the well on the left-hand side: need  $E < \inf\{j_\varphi(0) \mid K(\varphi) = 0, \varphi \neq 0\} = J(Q)$  (lowest mountain pass). Expect global existence in that case.

## Above the ground state energy

Theorem (Nakanishi-S. 2010)

Let  $E(u_0, u_1) < E(Q, 0) + \varepsilon^2$ ,  $(u_0, u_1) \in \mathcal{H}_{\text{rad}}$ . In  $t \geq 0$  for NLKG:

1. *finite time blowup*
2. *global existence and scattering to 0*
3. *global existence and scattering to  $Q$ :  $u(t) = Q + v(t) + o_{\mathcal{H}^1}(1)$  as  $t \rightarrow \infty$ , and  $\dot{u}(t) = \dot{v}(t) + o_{L^2}(1)$  as  $t \rightarrow \infty$ ,  $\square v + v = 0$ ,  $(v, \dot{v}) \in \mathcal{H}$ .*

*All 9 combinations of this trichotomy allowed as  $t \rightarrow \pm\infty$ .*

- Applies to  $\dim = 3$ ,  $|u|^{p-1}u$ ,  $7/3 < p < 5$ , or  $\dim = 1$ ,  $p > 5$ .
- Third alternative forms the **center stable manifold** associated with  $(\pm Q, 0)$ . Linearized operator  $L_+ = -\Delta + 1 - 3Q^2$  has spectrum  $\{-k^2\} \cup [1, \infty)$  on  $L_{\text{rad}}^2(\mathbb{R}^3)$ . **Gap**  $[0, 1)$  difficult to verify, Costin-Huang-S., 2011.
- $\exists$  1-dim. **stable, unstable manifolds** at  $(\pm Q, 0)$ . **Stable manifolds:** Duyckaerts-Merle, Duyckaerts-Holmer-Roudenko 2009

## The invariant manifolds




Figure : Stable, unstable, center-stable manifolds

## Hyperbolic dynamics near $\pm Q$

Linearized operator  $L_+ = -\Delta + 1 - 3Q^2$

- $\langle L_+ Q | Q \rangle = -2\|Q\|_4^4 < 0$
- $L_+ \rho = -k^2 \rho$  unique negative eigenvalue, no kernel over radial functions
- Gap property:  $L_+$  has no eigenvalues in  $(0, 1]$ , no threshold resonance (delicate!) Use Kenji Yajima's  $L^p$ -boundedness for wave operators.

Plug  $u = Q + v$  into cubic NLKG:

$$\ddot{v} + L_+ v = N(Q, v) = 3Qv^2 + v^3$$

Rewrite as a Hamiltonian system:

$$\partial_t \begin{pmatrix} v \\ \dot{v} \end{pmatrix} = \begin{bmatrix} 0 & 1 \\ -L_+ & 0 \end{bmatrix} \begin{pmatrix} v \\ \dot{v} \end{pmatrix} + \begin{pmatrix} 0 \\ N(Q, v) \end{pmatrix}$$

Then  $\text{spec}(A) = \{k, -k\} \cup i[1, \infty) \cup i(-\infty, -1]$  with  $\pm k$  simple evals. Formally:

$X_s = P_1 L^2$ ,  $X_u = P_{-1} L^2$ ,  $X_c$  is the rest.

## Spectrum of matrix Hamiltonian




Figure : Spectrum of nonselfadjoint linear operator in phase space

## Numerical 2-dim section through $\partial S_+$ (with R. Donninger)





Figure :  $(Q + Ae^{-r^2}, Be^{-r^2})$

- soliton at  $(A, B) = (0, 0)$ ,  $(A, B)$  vary in  $[-9, 2] \times [-9, 9]$
- **RED**: global existence, **WHITE**: finite time blowup, **GREEN**:  $\mathcal{PS}_+$ , **BLUE**:  $\mathcal{PS}_-$
- Our results apply to a neighborhood of  $(Q, 0)$ , boundary of the red region looks smooth (caution!)

## Variational structure above $E(Q, 0)$



- Solution can pass through the balls. Energy is no obstruction anymore as in the Payne-Sattinger case.
- **Key to description of the dynamics: One-pass (no return) theorem.** The trajectory can make only one pass through the balls.
- **Point: Stabilization of the sign of  $K(u(t))$ .**

## One-pass theorem (non-perturbative)




Figure : Possible returning trajectories

Such trajectories are **excluded** by means of an indirect argument using a variant of the **virial argument** that was essential to the **rigidity step of concentration compactness**.

## One-pass theorem

**Crucial no-return property:** Trajectory does **not return to balls around**  $(\pm Q, 0)$ . Suppose it did; Use *virial identity*

$$\partial_t \langle w \dot{u} | Au \rangle = - \int_{\mathbb{R}^3} (|\nabla u|^2 - \frac{3}{4}|u|^4) dx + \text{error}, \quad A = \frac{1}{2}(x\nabla + \nabla x)$$

where  $w = w(t, x)$  is a **space-time cutoff** that lives on a **rhombus**, and the “error” is controlled by the **external energy**.

Finite propagation speed  $\Rightarrow$  error controlled by **free energy outside large balls** at times  $T_1, T_2$ .

Integrating between  $T_1, T_2$  gives **contradiction**; the **bulk** of the integral of  $K_2(u(t))$  here comes from **exponential ejection** mechanism near  $(\pm Q, 0)$ .

**Non-perturbative argument.**




Figure : Space-time cutoff for the virial identity

## Open problem

**Complete description of possible long-term dynamics:** Given focusing NLKG3 in  $\mathbb{R}^3$  with radial energy data, show that the solution either

- blows up in finite time
- exists globally, scatters to one of the stationary solutions  $-\Delta\varphi + \varphi = \varphi^3$  (including 0)

Moreover, describe dynamics, center-stable manifolds associated with  $\varphi$ .

**Evidence:** With dissipation given by  $-\alpha\partial_t u$  term, result holds (Burq-Raugel-S.).

**Critical equation:**  $\square u = u^5$  in  $\mathbb{R}^3$ , Duyckaerts-Kenig-Merle proved analogous result with rescaled ground-state profiles  $\sqrt{\lambda}W(\lambda x)$ ,  $W(x) = (1 + |x|^2/3)^{-\frac{1}{2}}$ .

**Obstruction:** Exterior energy estimates in DKM scheme fail in the KG case due to speed of propagation  $< 1$ .

## Center-stable manifold, $u^5$ critical equation

Nakanishi-S theorem applies to **nonradial NLKG, NLS**, different **subcritical** powers and dimensions. **Critical equations** exhibit similar, yet **qualitatively essentially different** phenomena due to **scaling symmetry**.

$$\ddot{u} - \Delta u = |u|^{2^*-2}u, \quad u(t, x) : \mathbb{R}^{1+d} \rightarrow \mathbb{R}, \quad 2^* = \frac{2d}{d-2},$$

Static **Aubin, Talenti solutions**

$$W_\lambda = T_\lambda W, \quad W(x) = \left[ 1 + \frac{|x|^2}{d(d-2)} \right]^{1-\frac{d}{2}},$$

$T_\lambda$  is  $\dot{H}^1$  preserving dilation

$$T_\lambda \varphi = \lambda^{d/2-1} \varphi(\lambda x)$$

**Positive radial solutions**  $-\Delta W - |W|^{2^*-2}W = 0$ . **Functionals:**

$$J(\varphi) := \int_{\mathbb{R}^d} \left[ \frac{1}{2} |\nabla \varphi|^2 - \frac{1}{2^*} |\varphi|^{2^*} \right] dx, \quad K(\varphi) := \int_{\mathbb{R}^d} [|\nabla \varphi|^2 - |\varphi|^{2^*}] dx$$

## Critical wave equation

Radial  $\dot{H}^1 \times L^2$ ,  $E(\vec{\varphi}) < J(W) + \varepsilon^2$ , outside soliton tube

$$\mathcal{S} = \{\pm \vec{W}_\lambda \mid \lambda > 0\} + O(\varepsilon)$$

There exists four open disjoint sets which lead to all combinations of FTB/GE and scattering to 0 as  $t \rightarrow \pm l$ .

- Krieger-Nakanishi-S. 2013: complete description of all solutions with energy  $E(\vec{\varphi}) < J(W) + \varepsilon^2$ . Type-I conjecture!
- center-stable manifold exists in  $\dot{H}^1 \times L^2$ , contains all  $W_\lambda$ , solutions with  $\lambda \rightarrow 0, \infty$  (but Krieger-S. 05 showed that in a stronger non-invariant topology exists codim-1 manifold with global solutions,  $\lambda(t) \rightarrow \lambda_* \in (0, \infty)$ ).
- Inside the soliton tube there exist blowup solutions, as found by Krieger-S.-Tataru 06. Then Duyckaerts-Kenig-Merle 09 showed that all type II blowup are of the KST form, as long as energy below  $2J(Q)$ . So trapping by the soliton tube cannot mean scattering to  $\{W_\lambda\}$  as it did in the subcritical case.

## Equivariant wave maps

$u : \mathbb{R}_{t,x}^{1+2} \rightarrow \mathbb{S}^2$  satisfies **WM equation**

$$\square u \perp T_u \mathbb{S}^2 \Leftrightarrow \square u = u(|\partial_t u|^2 - |\nabla u|^2)$$

as well as **equivariance assumption**  $u \circ R = R \circ u$  for all  $R \in SO(2)$




Figure : Equivariance and Riemann sphere

## Equivariant wave maps

$u(t, r, \phi) = (\psi(t, r), \phi)$ , spherical coordinates,  $\psi$  angle from north pole satisfies

$$\psi_{tt} - \psi_{rr} - \frac{1}{r}\psi_r + \frac{\sin(2\psi)}{2r^2} = 0, \quad (\psi, \psi_t)(0) = (\psi_0, \psi_1)$$

- **Conserved energy**

$$E(\psi, \psi_t) = \int_0^\infty \left( \psi_t^2 + \psi_r^2 + \frac{\sin^2(\psi)}{r^2} \right) r \, dr$$

- $\psi(t, \infty) = n\pi, n \in \mathbb{Z}$ , **homotopy class = degree =  $n$**
- **stationary solutions = harmonic maps =  $0, \pm Q(r/\lambda)$** , where  $Q(r) = 2 \arctan r$ . This is the identity  $\mathbb{S}^2 \rightarrow \mathbb{S}^2$  with **stereographic projection** onto  $\mathbb{R}^2$  as domain (**conformal map!**).

## Large data results for equivariant wave maps

Theorem (Côte, Kenig, Lawrie, S. 2012)

Let  $(\psi_0, \psi_1)$  be smooth data.

1. Let  $E(\psi_0, \psi_1) < 2E(Q, 0)$ , degree 0. Then the solution exists globally, and scatters (energy on compact sets vanishes as  $t \rightarrow \infty$ ). For any  $\delta > 0$  there exist data of energy  $< 2E(Q, 0) + \delta$  which blow up in finite time.
2. Let  $E(\psi_0, \psi_1) < 3E(Q, 0)$ , degree 1. If the solution  $\psi(t)$  blows up at time  $t = 1$ , then there exists a continuous function,  $\lambda : [0, 1) \rightarrow (0, \infty)$  with  $\lambda(t) = o(1 - t)$ , a map  $\vec{\varphi} = (\varphi_0, \varphi_1) \in \mathcal{H}$  with  $E(\vec{\varphi}) = E(\vec{\psi}) - E(Q, 0)$ , and a decomposition


$$\vec{\psi}(t) = \vec{\varphi} + (Q(\cdot/\lambda(t)), 0) + \vec{\epsilon}(t) \quad (\star)$$

s.t.  $\vec{\epsilon}(t) \in \mathcal{H}$ ,  $\vec{\epsilon}(t) \rightarrow 0$  in  $\mathcal{H}$  as  $t \rightarrow 1$ .

## Large data results for equivariant wave maps


- For **degree 1** have an analogous classification to (★) for **global solutions**.
- Côte 2013: **bubble-tree** classification for **all** energies **along a sequence** of times.  
**Open problems:** (A) all times, rather than a sequence (B) construction of bubble trees.
- Duyckaerts, Kenig, Merle 12 established classification results for  $\square u = u^5$  in  $\dot{H}^1 \times L^2(\mathbb{R}^3)$  with  $W(x) = (1 + |x|^2/3)^{-\frac{1}{2}}$  instead of  $Q$ .
- Construction of (★) by Krieger-S.-Tataru 06 in finite time, Donninger-Krieger 13 in *infinite time* (for critical NLW)
- **Crucial role is played by Michael Struwe's bubbling off theorem (equivariant):** if blowup happens, then there exists a sequence of times approaching blowup time, such that a rescaled version of the wave map approaches locally in energy space a harmonic map of positive energy.

## Struwe's cuspidal energy concentration



Rescalings converge in  $L^2_{t,r}$ -sense to a **stationary wave map** of positive energy, i.e., a **harmonic map**.

## Asymptotic exterior energy



$\square u = 0$ ,  $u(0) = f \in \dot{H}^1(\mathbb{R}^d)$ ,  $u_t(0) = g \in L^2(\mathbb{R}^d)$  radial

Duyckaerts-Kenig-Merle 2011: for all  $t \geq 0$  or  $t \leq 0$  have  $E_{\text{ext}}(\vec{u}(t)) \geq cE(f, g)$   
provided dimension odd.  $c > 0$ ,  $c = \frac{1}{2}$

Heuristics: incoming vs. outgoing data.

## Exterior energy: even dimensions

Côte-Kenig-S. 2012: This **fails in even dimensions**.

$d = 2, 6, 10, \dots$  holds for data  $(0, g)$  but fails in general for  $(f, 0)$ .

$d = 4, 8, 12, \dots$  holds for data  $(f, 0)$  but fails in general for  $(0, g)$ .

Fourier representation, Bessel transform, dimension  $d$  reflected in the phase of the Bessel asymptotics, computation of the asymptotic exterior energy as  $t \rightarrow \pm\infty$ .

For our  $3E(Q, 0)$  theorem we need  $d = 4$  result; rather than  $d = 2$  due to repulsive  $\frac{\psi}{r^2}$ -potential coming from  $\frac{\sin(2\psi)}{2r^2}$ .

$(f, 0)$  result suffices by Christodoulou, Tahvildar-Zadeh, Shatah results from mid 1990s. Showed that at blowup  $t = T = 1$  have vanishing kinetic energy

$$\lim_{t \rightarrow 1} \frac{1}{1-t} \int_t^1 \int_0^{1-t} |\dot{\psi}(t, r)|^2 r dr dt = 0$$

No result for Yang-Mills since it corresponds to  $d = 6$

## Exterior energy: odd dimensions

Duyckaerts-Kenig-Merle: in radial  $\mathbb{R}^3$  one has for all  $R \geq 0$

$$\max_{\pm} \lim_{t \rightarrow \pm\infty} \int_{|x| > t+R} |\nabla_{t,x} u|^2 dr \geq c \int_{|x| > R} [(ru)_r^2 + (ru)_t^2] dr$$

**Note:** RHS is **not** standard energy! **Orthogonal projection** perpendicular to **Newton potential**  $(r^{-1}, 0)$  in  $H^1 \times L^2(\mathbb{R}^3 : r > R)$ .

Kenig-Lawrie-S. 13 noted this projection and extended the exterior energy estimate to  $d = 5$ : project perpendicular to plane  $(\xi r^{-3}, \eta r^{-3})$  in  $H^1 \times L^2(\mathbb{R}^5 : r > R)$

Kenig-Lawrie-Liu-S. 14 **all odd dimensions**, projections off of similar but larger and more complicated linear subspaces.

**Relevance:** Exterior wave maps in  $\mathbb{R}^3$  with arbitrary degree of equivariance lead to **all odd dimensions**.

## Exterior wave maps

Consider equivariant wave maps from  $\mathbb{R}^3 \setminus B(0, 1) \rightarrow \mathbb{S}^3$  with **Dirichlet condition** at  $R = 1$ . **Supercritical becomes subcritical**, easy to obtain global smooth solutions.

**Conjecture by Bizon-Chmaj-Maliborski 2011:** All smooth solutions scatter to the unique harmonic map in their degree class.

### Results:

- **Lawrie-S. 2012:** Proved for degree 0 and **asymptotic stability** for degree 1. Follows Kenig-Merle concentration compactness approach with rigidity argument carried out by a virial identity (complicated).
- **Kenig-Lawrie-S. 2013:** Proved for all degrees in equivariance class 1. Uses exterior energy estimates instead of virial.
- **Kenig-Lawrie-Liu-S. 2014:** Proved for all degrees and all equivariance classes. Requires exterior energy estimates in all odd dimensions.

**Soliton resolution conjecture** holds in this case.

THANK YOU FOR YOUR ATTENTION!